



## Effects of a Self-Management Short Course Instruction on Glycemic Control in Adults with Diabetes Mellitus

Mohammad Afshar<sup>1</sup>, Neda Mirbagher Ajorpaz<sup>1\*</sup>

<sup>1</sup>Trauma Nursing Research Center, Kashan University of Medical Sciences, Kashan, IR Iran

### ARTICLE INFO

**Article type:**  
Original Article

**Article history:**  
Received: 12 May 2012  
Revised: 13 Jun 2012  
Accepted: 08 Jul 2012

**Keywords:**  
Teaching  
Self Care  
Diabetes Mellitus

### ABSTRACT

**Background:** Education is an integral part of the treatment in diabetes mellitus. Attendance at long courses might not be convenient for many patients.

**Objectives:** The current study was conducted to evaluate the effectiveness of self-management, short course instruction on glycemic control in adults with diabetes mellitus.

**Patients and Methods:** A total of 60 patients with diabetes mellitus were randomly allocated into intervention (n = 30) and control (n = 30) groups. Fasting blood sugar (FBS) and blood sugar (BS) (5pm) tests were conducted. The intervention group received instruction about self-management in diabetes mellitus for two hours, during two sessions. They were followed-up for three months with telephone calls. Patients asked any questions they had during these calls. After three months the patients' FBS and BS were recorded again. The same process took place in the control group without training. Independent sample t-test and chi-square tests were used to analyze data using SPSS version 16.0.

**Results:** The sample included 60 patients with a mean age of  $46 \pm 2.14$  years. The FBS dropped from 151 mg/dL to 110 mg/dL in the intervention group ( $P = 0.02$ ). While it increased from 146 mg/dL to 150 mg/dL in the control group. The BS also decreased from 231 mg/dL to 196 in the intervention group. ( $P = 0.05$ ), but it increased from 240 to 247 in the control group ( $P = 0.09$ ). There was a significant difference in FBS and BS tests in the two groups after three months. ( $P = 0.002, P = 0.05$ ), respectively.

**Conclusions:** The results showed that a short course of instruction is effective in glycemic control. It is suggested that further research is conducted to evaluate the effectiveness of self-management long course instruction on glycemic control in adults with diabetes mellitus.

Published by Kowsar Corp, 2012. cc 3.0.

### ► Implication for health policy/practice/research/medical education:

Health authorities are responsible for integrating self-management instruction and follow-up programs in the course of diabetes care.

### ► Please cite this paper as:

Afshar M, Mirbagher Ajorpaz N. Effects of a Self-Management Short Course Instruction on Glycemic Control in Adults with Diabetes Mellitus. *Nurs Midwifery Stud.* 2012;1(1): 7-11. DOI: 10.5812/nms.7892

## 1. Background

Diabetes mellitus is a relatively common chronic disease with no cure at present (1). Type 1 diabetes usually develops in childhood and adolescence, whereas type

2 diabetes, is not common before the age of 40 years (2, 3). There is evidence that the incidence of diabetes mellitus is rising in the world (4, 5). The International Diabetes Federation estimates that in 2010, over 200 million

\* Corresponding author: Neda Mirbagher Ajorpaz, Trauma Nursing Research Center, Kashan University of Medical Sciences, Kashan, IR Iran. Tel: +98-3615550021, Fax: +98-3615556633, E-mail: mirbagher\_n@kaumc.ac.ir

people in the world were affected by diabetes (6). This encompasses approximately 6% of the world's population (7). People with diabetes have elevated risks for; retinopathy, renal failure, neuropathy, atherosclerosis, peripheral artery disease, coronary heart disease and cerebrovascular disease. It is expected that better diabetes control will postpone or even prevent some of these complications (8). Although diabetes cannot be cured, the disease can be managed by pharmacological and non-pharmacological strategies (9). Patients' understanding of diabetes mellitus and its treatment have been viewed as essential to the management of this complex, chronic illness. To this end, the American Diabetes Association and other national agencies have recommended diabetic patients' education as an integral part of their treatment (10, 11). Recent discussions about educational interventions for patients with diabetes have focused on the potential benefits of simultaneously improving patients' understanding of diabetes, providing support for healthy behavior changes, and empowering patients (12, 13). Formal diabetes education has often focused on lifestyle modifications such as; dietary change, exercise, and self-measurement of blood glucose. Educational interventions have been shown to increase patients' knowledge of diabetes and self-care activities (14), especially in the short-term, but the results of these interventions on important long-term health outcomes remain unclear (12-15). Improving patients' knowledge of diabetes through educational efforts could lead more patients to take an active role in obtaining the necessary preventive care (16). It is widely recommended that educational interventions should be an integral part of diabetes care (17, 18). Educational interventions primarily teach diabetes-related knowledge and skills required for self-management, including correctly testing blood-glucose levels and injecting insulin (5). A multidisciplinary educational program of at least six to 25 hours is recommended for diabetes education (19). However, many patients are not able to attend these programs and long courses might be exhausting or inconvenient for some patients. In two similar reports, 76-85% of diabetic patients had poor knowledge and 33.3% had poor performance on self-care, despite routine patient education programs that had been delivered in diabetes centers. Consequently, the need for improved patient education programs and identifying ways to empower patients in diabetic care has been emphasized (14).

## 2. Objectives

The aim of the current study was to evaluate the effects of self-management, short course instruction on glycemic control in adults with diabetes mellitus.

## 3. Patients and Methods

A randomized controlled pre- and post-test design was employed to verify the effect of self-management, short

course instruction on glycemic control in patients with diabetes mellitus attending the outpatient clinic of the Shahid Beheshti Hospital, Kashan, Iran. The inclusion criteria were; age between 20-65 years, good general health, history of diabetes and injecting insulin for at least one year. Patients who were pregnant or diagnosed with end-stage renal disease or any other severe condition were excluded from the study. Exclusion criteria were; failure to follow the education program. A sample of 60 patients participated in the study. The purpose of the study was explained and informed consent was obtained. Numbers for the 60 patients were listed and using a random number's table, the names were allocated into two equal groups. Initially, the two groups completed demographic questionnaires (age, sex, race, education, income, and marital status) and patient clinical characteristics (self-reported health status, number of co-morbidities and years with diabetes). Then, fasting blood sugar (FBS) and blood sugar (BS) (5 pm) tests were carried out in both of the two groups. In the intervention group, face-to-face instruction was given by a researcher. The content of the instruction were lifestyle modifications such as; dietary changes, exercise, self-measurement of blood glucose, and blood pressure control. The instructions were carried out in two sessions; the duration of each session was one hour, with a one-week interval. An educational booklet was also given to the intervention group. Patients were followed for three months by telephone, and they could also ask questions. After three months, FBS and BS tests were taken and recorded in terms of mg/dL. The same process took place in the control group which did not receive any training. The mean score of FBS-BS was then calculated for each participant. t-test and chi-square were used to analyze data using SPSS version 16.0. A P value less than 0.05 was considered to be significant. This study received a grant from the Institutional Review Board (IRB) and the study was approved by the Kashan University of Medical Sciences (KAUMS), ethics approval was received from the Ethics Committee of KAUMS.

## 4. Results

The study included 60 patients with a mean age of  $46 \pm 2.14$  years. Other characteristics of the participants are shown in *Table 1*. The intervention and control groups showed no significant differences in age, sex, duration of diabetes, and marital status (*Table 1*). Following the intervention, the mean score of FBS dropped from 151 mg/dL to 110 mg/dL in the intervention group ( $P = 0.02$ ), while the mean score of the FBS increased from 146 mg/dL to 150 mg/dL in the control group. The mean score of the BS also decreased from 231 mg/dL to 196 in the intervention group, and it showed an increase from 240 to 247 in the control group (*Table 2*). There was a significant difference in FBS and BS tests in the groups three months after the education sessions. ( $P = 0.002, P = 0.05$ ) (*Table 3*).

**Table 1.** Demographic Characteristics of Patients

|                                  | Intervention Group, No. (%) | Control Group, No. (%) | X <sup>2</sup> | P value |
|----------------------------------|-----------------------------|------------------------|----------------|---------|
| <b>Age, y</b>                    |                             |                        | 1.34           | 0.2     |
| 20 - 30                          | 2 (6.6)                     | 4 (13.3)               |                |         |
| 30 - 40                          | 8 (26.6)                    | 9 (30)                 |                |         |
| 40 - 50                          | 15 (50)                     | 12 (40)                |                |         |
| 50 - 60                          | 5 (16.6)                    | 5 (16.6)               |                |         |
| <b>Gender</b>                    |                             |                        | 2.01           | 0.32    |
| Female                           | 18 (60)                     | 21 (70)                |                |         |
| Male                             | 12 (40)                     | 9 (30)                 |                |         |
| <b>Marital Status</b>            |                             |                        | 1.74           | 0.1     |
| Single                           | 7 (23.3)                    | 1 (3.3)                |                |         |
| Married                          | 20 (66.6)                   | 24 (80)                |                |         |
| Widow                            | 3 (10)                      | 5 (16.6)               |                |         |
| <b>Duration with diabetes, y</b> |                             |                        | 2.13           | 0.84    |
| 1 - 5                            | 17 (56.6)                   | 15 (50)                |                |         |
| 5 - 10                           | 8 (26.6)                    | 9 (30)                 |                |         |
| Up to 10                         | 5 (16.6)                    | 6 (20)                 |                |         |
| <b>Insulin use in day</b>        |                             |                        | 1.76           | 0.15    |
| Once                             | 18 (60)                     | 15 (50)                |                |         |
| Twice                            | 12 (40)                     | 15 (50)                |                |         |

**Table 2.** Fasting Blood Sugar and Blood Sugar Level Before and After Instruction in the Two Groups

|                  | Intervention Group |                  |         |        |
|------------------|--------------------|------------------|---------|--------|
|                  | Before, Mean ± SD  | After, Mean ± SD | P value | t-Test |
| <b>FBS</b>       | 151 ± 2.4          | 110 ± 2.1        | 0.02    | 3.45   |
| <b>BS (5 pm)</b> | 231 ± 2.5          | 196 ± 2.8        | 0.05    | 3.01   |
|                  | Control Group      |                  |         |        |
|                  | Before, Mean ± SD  | After, Mean ± SD | P value | t-Test |
| <b>FBS</b>       | 146 ± 1.3          | 150 ± 2.2        | 0.14    | 1.78   |
| <b>BS (5 pm)</b> | 240 ± 2.8          | 247 ± 3.2        | 0.09    | 2.65   |

Abbreviations: FBS, fasting blood sugar; BS, blood sugar

**Table 3.** Comparison of FBS and BS Tests After Instruction in the Two Groups

|           | Intervention Group, Mean ± SD | Control Group, Mean ± SD | P value | t-Test |
|-----------|-------------------------------|--------------------------|---------|--------|
| FBS       | 110 ± 2.1                     | 150 ± 2.2                | 0.002   | 4.65   |
| BS (5 pm) | 196 ± 2.8                     | 247 ± 3.2                | 0.05    | 3.98   |

Abbreviations: FBS, fasting blood sugar; BS, blood sugar

## 5. Discussion

In the present study, the effect of a short-course instruction on self-management in diabetes mellitus was examined. The results indicated that a short course on self-management instruction had a significant effect on glycemic control in diabetes mellitus. Education of diabetic patients often focuses on self-management activities, including; diet, exercise, self-measurement of blood glucose, problem-solving skills, and methods for coping

with diabetes (20-22). Norris *et al.* showed that patients who received diabetes education from a clinician or a more knowledgeable person, were more likely to perform self-management activities (23). The current study supported the results of studies conducted by Norris *et al.* (23) and Persell *et al.* (24) which showed that changes in blood-glucose levels before and after instruction were significant. Gage *et al.* also showed that educational and psychosocial programs are effective in diabetes control

(4). In a study performed by Keers et al. (2004), it was pointed out that there was a meaningful difference in FBS and BS levels before and after giving instruction (25). Malone et al. randomized patients presenting with severe diabetic complications into those receiving care education and those with no education. After two years of follow-up, diabetic complication rates were three times lower in the intervention than in the control group (26). However, Jeffcoate et al. conducted a similar study design, and failed to verify these findings (27). In another study, there were no significant differences shown between intervention and control groups after a short-course of instruction on self-management in diabetes mellitus (28). The World Health Organization (WHO), in two separate reports declared that diabetes is increasing in developing countries, and invited members to investigate the importance of diabetes in their individual countries and requested them to apply appropriate preventive interventions to eliminate the complications of diabetes (29). The present results show that short-course teaching programs using self-management methods, which are accompanied by follow-up, enable patients to increase control of their blood sugar. A short-course of instruction is easier to attend, and it can provide a normal life for diabetic patients, not to mention the time and money that can also be saved. However, these types of interventions require professional trainers. The long-term effects of short-term instructions on glycemic control, quality of life and self-efficacy need to be further evaluated and the identification of patients who might benefit from this method of instruction also need to be determined.

## Acknowledgments

Our thanks go to all the patients who participated in this research.

## Authors' Contribution

Mohammad Afshar was responsible for study conception and design, data collection and participated in preparing the first draft. Neda Mirbagher Ajorpaz prepared the first draft of the manuscript, performed the data analysis, and made revisions to the paper.

## Financial Disclosure

None declared.

## Funding/Support

This study was funded and supported by the Deputy of Research, Kashan University of Medical Sciences (KAUMS), Grant No: 8608.

## References

1. Grau T, Almazan Arjona JA, Luna A, Chamorro Quiros J, Lord Rodriguez T, Casimiro C, et al. [Evaluation of palatability of two special oral diets for institutionalized elderly diabetics, Glucerna SR vs Resource Diabet]. *Nutr Hosp*. 2004;**19**(5):292-9.
2. Masding M, Jones J, Bartley E, Sandeman D. Assessment of blood pressure in patients with Type 2 diabetes: comparison between home blood pressure monitoring, clinic blood pressure measurement and 24-h ambulatory blood pressure monitoring. *Diabetic Med*. 2001;**18**(6):431-7.
3. Tudor C, Bacanu G, Serban V, Staicu M, Tudor R, Negrisanu G. [The early detection of diabetes mellitus and the active dispensary care of 1200 subjects with a genetic predisposition]. *Med Interna*. 1991;**43**(1-2):141-9.
4. Gage H, Hampson S, Skinner TC, Hart J, Storey L, Foxcroft D, et al. Educational and psychosocial programmes for adolescents with diabetes: approaches, outcomes and cost-effectiveness. *Patient Educ Couns*. 2004;**53**(3):333-46.
5. Sasaki H, Kawasaki T, Ogaki T, Kobayashi S, Itoh K, Yoshimizu Y, et al. The prevalence of diabetes mellitus and impaired fasting glucose/glycaemia (IFG) in suburban and rural Nepal-the communities-based cross-sectional study during the democratic movements in 1990. *Diabetes Res Clin Pract*. 2005;**67**(2):167-74.
6. Schipf S, Werner A, Tamayo T, Holle R, Schunk M, Maier W, et al. Regional differences in the prevalence of known Type 2 diabetes mellitus in 45-74 years old individuals: results from six population-based studies in Germany (DIAB-CORE Consortium). *Diabet Med*. 2012;**29**(7):e88-95.
7. Ruusunen A, Voutilainen S, Karhunen L, Lehto S, Tolmunen T, Keinänen-Kiukaanniemi S, et al. How does lifestyle intervention affect depressive symptoms? Results from the Finnish Diabetes Prevention Study. *Diabetic Med*. 2012.
8. Cafazzo JA, Casselman M, Hamming N, Katzman DK, Palmert MR. Design of an mHealth app for the self-management of adolescent type 1 diabetes: a pilot study. *J Med Internet Res*. 2012;**14**(3):e70.
9. Minet L, Moller S, Vach W, Wagner L, Henriksen JE. Mediating the effect of self-care management intervention in type 2 diabetes: a meta-analysis of 47 randomised controlled trials. *Patient Educ Couns*. 2010;**80**(1):29-41.
10. American Diabetes A. Standards of medical care for patients with diabetes mellitus. *Diabetes Care*. 2003;**26** (Suppl 1):S33-50.
11. Funnell MM, Brown TL, Childs BP, Haas LB, Hoseney GM, Jensen B, et al. National standards for diabetes self-management education. *Diabetes Care*. 2010;**33** (Suppl 1):S89-96.
12. Glasgow RE, Anderson RM. In diabetes care, moving from compliance to adherence is not enough. Something entirely different is needed. *Diabetes Care*. 1999;**22**(12):2090-2.
13. Glasgow RE, Toobert DJ, Riddle M, Donnelly J, Mitchell DL, Calder D. Diabetes-specific social learning variables and self-care behaviors among persons with type II diabetes. *Health Psychol*. 1989;**8**(3):285-303.
14. Adib-Hajbaghery M, Alinaqipoor T. Comparing the Effects of Two Teaching Methods on Healing of Diabetic Foot Ulcer. *J Caring Sci*. 2012;**1**(1):17-24.
15. Coffman M, Norton C, Beene L. Diabetes symptoms, health literacy, and health care use in adult Latinos with diabetes risk factors. *J Cult Divers*. 2012;**19**(1):4.
16. Gumbs J. Relationship between diabetes self-management education and self-care behaviors among African American women with type 2 diabetes. *J Cult Divers*. 2012;**19**(1):18.
17. Morrison S, Dashiff C, Abdullatif H, Moreland E. Parental separation anxiety and diabetes self-management of older adolescents: a pilot study. *Pediatr Nurs*. 2012;**38**(2):88-95.
18. Rodbard HW, Bays HE, Gavin JR, 3rd, Green AJ, Bazata DD, Lewis SJ, et al. Rate and risk predictors for development of self-reported type-2 diabetes mellitus over a 5-year period: the SHIELD study. *Int J Clin Pract*. 2012;**66**(7):684-91.
19. Mensing C, Boucher J, Cypress M, Weinger K, Mulcahy K, Barta P, et al. National standards for diabetes self-management education. *Diabetes Care*. 2002;**25** (Suppl 1):S149-56.
20. Didarloo A, Shojaeizadeh D, Gharaaghaji A, Habibzadeh H, Sh N, Pourali R. Prediction of Self-Management Behavior among Iranian Women with Type 2 Diabetes: Application of the Theory of Reasoned Action along with Self-Efficacy (ETRA). *Iran Red Crescent Med J*. 2012;**14**(2):85-94.
21. Hartayu TS, Mi MI, Suryawati S. Improving of Type 2 Diabetic Pa-

- tients' Knowledge, Attitude and Practice Towards Diabetes Self-care by Implementing Community-Based Interactive Approach-Diabetes Mellitus Strategy. *BMC Res Notes*. 2012;**5**:315.
22. Karter AJ, Ackerson LM, Darbinian JA, D'Agostino RB, Jr., Ferrara A, Liu J, et al. Self-monitoring of blood glucose levels and glycemic control: the Northern California Kaiser Permanente Diabetes registry. *Am J Med*. 2001;**111**(1):1-9.
  23. Norris SL, Engelgau MM, Narayan KM. Effectiveness of self-management training in type 2 diabetes: a systematic review of randomized controlled trials. *Diabetes Care*. 2001;**24**(3):561-87.
  24. Persell SD, Keating NL, Landrum MB, Landon BE, Ayanian JZ, Borbas C, et al. Relationship of diabetes-specific knowledge to self-management activities, ambulatory preventive care, and metabolic outcomes. *Prev Med*. 2004;**39**(4):746-52.
  25. Keers JC, Blaauwwekel EE, Hania M, Bouma J, Scholten-Jaegers SM, Sanderman R, et al. Diabetes rehabilitation: development and first results of a Multidisciplinary Intensive Education Program for patients with prolonged self-management difficulties. *Patient Educ Couns*. 2004;**52**(2):151-7.
  26. Malone JM, Snyder M, Anderson G, Bernhard VM, Holloway GA, Jr., Bunt TJ. Prevention of amputation by diabetic education. *Am J Surg*. 1989;**158**(6):520-3; discussion 3-4.
  27. Jeffcoate W, Radford K, Ince P, Smith M. Randomized controlled trial of education in the prevention of foot ulcer recurrence in diabetes. *Diabetologia*. 2007;**50** (Suppl 1):S457.
  28. Ali M, Schifano F, Robinson P, Phillips G, Doherty L, Melnick P, et al. Impact of community pharmacy diabetes monitoring and education programme on diabetes management: a randomized controlled study. *Diabet Med*. 2012;**29**(9):e326-33.
  29. Alwan A. Global status report on noncommunicable diseases 2010. WHO; 2011.