



Effects of simulation-based education on Ghanaian parents' knowledge, attitude and management of fever control

Batool Pouraboli ¹, Aida Safaiee Fakhri ², Jamalodin Begjani ¹, Abdulai Mohammed Hardi ^{3*}, Hadi Ranjbar ⁴, Mohammad Eghbal Heidari ⁵

¹ Department of Pediatric and Neonatal Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

² School of Public Health and Paraclinics, Qazvin University of Medical Sciences, Qazvin, Iran

³ Pediatric Nursing and Neonatal Intensive Care Unit Department, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

⁴ Mental Health Research Center, Psychosocial Health Research Institute, Iran University of Medical Sciences, Tehran, Iran

⁵ Student Scientific Research Center, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

* **Corresponding author:** Abdulai Mohammed Hardi, School of Nursing and Midwifery, Pediatric Nursing and Neonatal Intensive Care Unit Department, Tehran University of Medical Sciences, Tehran, Iran. E-mail: hajjmhadi@gmail.com

Received: 25 December 2022 **Revised:** 1 August 2023 **Accepted:** 7 August 2023 **e-Published:** 22 September 2023

Abstract

Background: Childhood fever is very discomforting and most parents are usually anxious when children exhibit fever, leading to unnecessary visits and admissions into pediatric emergency centers.

Objectives: This study aimed to determine the effects of simulation-based education on parents' knowledge, attitude, and management of fever in children.

Methods: A non-randomized quasi-experimental study using convenient sampling was conducted to recruit parents of children aged 3 months to 8 years admitted to the Tamale Teaching Hospital in Tamale, Ghana. The study was conducted from July to September 2019. Eighty parents were equally assigned into an intervention group and a control group and respectively educated using simulation-based and routine education approaches on childhood fever. Data were collected using a four-part questionnaire and analyzed using descriptive statistics, Chi-square, independent samples t, and paired t tests as well as analysis of covariance.

Results: Except for the mean baseline knowledge ($P < 0.0001$), there were no significant differences between the two groups in mean pretest scores. The mean scores of fever knowledge, attitude, and management of parents in the simulation-based education group were 21.63 ± 2.40 , 28.48 ± 6.03 , and 25.15 ± 4.23 at baseline and increased to 26.38 ± 4.36 , 28.48 ± 6.03 , and 32.33 ± 5.18 , respectively ($P < 0.0001$). All mean scores on the posttest were better than on the pretest in both groups. However, at the posttest, mean scores for knowledge, attitude, and management were significantly higher in the simulation-based education than in the routine education group ($P < 0.05$).

Conclusion: Compared with routine education, simulation-based education was more effective in improving parents' knowledge, attitude, and management of fever. We therefore, recommend the use of similar simulation-based methods in educating parents regarding caring for acute fever in their children.

Keywords: Fever, Education, Knowledge, Attitude, Parents, Child.

Introduction

Fever is defined as a body temperature above 38°C and usually results from injuries, infections, and immune responses.^[1,2] It is the main reason for up to 70% of pediatric emergencies.^[3] Fever is usually benign and useful in producing antibodies, and usually resolves without complications, except in severe bacterial infections.^[4] However, parents are anxious when children have fever.^[5]

About 32% to 56% of children with fever often come to health facilities for management,^[6] and about 4% experience febrile convulsions.^[7] Some studies in Africa have shown that most parents self-diagnose and self-medicate for fever in their children. They usually use physical methods such as sponging with cold water, herbal medicines (grounded garlic), and homeopathic medicines to treat fever.^[8,9] The revised Integrated Management of

Childhood Illnesses Strategy is the protocol used to manage all fever-related childhood illnesses.^[10,11] Despite the existence of the protocol, parents and healthcare workers still express fears about managing fever.^[12] Innovative care approaches and parental education on childhood fever are therefore necessary.^[1,13]

Simulation-based education methods mimic and amplify real-life experiences with guided experiences that replicate substantial aspects of the real world in interactive environments.^[14] Nurses can use simulation-based education in the form of experiments, scenarios, role playing/modeling, standardized patients, high human fidelity, videos, and other technological innovations, to educate their peers, clients, and students.^[15]

While traditional face-to-face education methods are still useful, simulation methods offer additional benefits by improving parents' information, motivation, and behavioral skills.^[16-20]

Many parents still lack appropriate knowledge, attitudes, and skills in managing fever in children.^[3,21] As a result, several studies have sought the best educational methods that impact knowledge, attitude, and practice of fever management.^[21] Consistently, positive outcomes have been revealed with patient satisfaction, improved fever knowledge, and positive attitudes even though the forms of education used varied.^[20,22-24] In a study, fever awareness increased by 68% in the group that was exposed to educational video messages and by 25% in the group that received a standard pamphlet.^[25] In another study, caregivers who had received a fever management video demonstrated better knowledge and attitude about fever in children than the group who watched a video about child safety.^[26] Another study by Moon *et al.*,^[27] showed that parents' knowledge improved better after watching short video messages than those who received text messages.^[28] In all of these studies video education was compared with methods such as educational texts or control videos but was not combined video education with face-to-face interactive methods. Given the lack of adherence to the Integrated Management of Childhood Illness's recommendations about childhood fever, it is necessary to compare the effectiveness of simulation-based education versus routine education on parents' knowledge, attitude, and management of fever control.

Objectives

This study aimed to determine the effects of simulation-based education on parents' knowledge, attitude, and management of fever in children.

Methods

Study design and participants

We conducted a nonrandomized quasi-experimental study at Tamale Teaching Hospital in Ghana. The study was conducted from July to September 2019. The sample size was calculated using the results of a previous study where simulation-based education on fever management in children increased the parental management behavior score from 2.19 ± 1.52 to 3.12 ± 1.3 .^[20] Then, using the formula for the comparison of two means, and considering a power of 0.80, a type one error of 0.05, and a possible dropout of 10%, we estimated that at least 40 participants were needed for each group.

Parents whose children were at least 3 months to 8 years old and were admitted to the pediatric ward due to acute fever were eligible for the study. Parents who declined participation, whose children were diagnosed with chronic fever-related illnesses or triaged to receive emergency care, and who were members of the health care team were excluded. Eligible parents were consecutively enrolled from two pediatric wards at Tamale Teaching Hospital in Ghana. A total of 102 parents were screened for eligibility, of which 80 eligible ones who gave consent were enrolled and equally assigned to two groups.

Data collection Instruments

The study instrument was made by the researcher based on a literature review^[20,29-31] and consists of 13 items on demographics, 15 yes/no or multiple-choice questions on knowledge, four questions on attitude, and 11 single and multiple questions on fever management (some questions have sub-items). In the knowledge section, every correct answer is scored as 1 for single response and 1 to 4 for multiple response questions, and the total score for the knowledge can range between 0 and 30.

The four items in the attitude sections have 14 sub-items and are rated on Likert-type responses ranging between zero and four, so that the total score can vary between 0 and 56. The total score in the management section can range between 0 and 44.

Five academics from the Pediatric Nursing Department of School of Nursing and Midwifery at Tehran University of Medical Sciences and five nurses from the pediatric ward of Tamale Teaching Hospital and Tamale Nursing and Midwifery Training College in Ghana confirmed the content and face validity of the study instrument. After piloting the instrument with 25 parents, The K-R 21 of the knowledge and attitude scales were 0.71 and 0.59, respectively, and the Cronbach's alpha of the management scale was 0.61. The same questionnaire was used for the

pretest and posttest that were conducted at baseline and two weeks after the educational interventions.

Interventions

A 12-minute video was developed by the research team and was viewed by parents in the intervention group. The video depicted a home setting with a mother narrating to her colleagues, her experiences and new fever information she had acquired during a recent visit to a health center. Four scenarios about fever control were presented verbally to help parents to identify and make appropriate decisions regarding likely future events, and lastly, a section of the video demonstrated the use of a thermometer. The video was paused after each section for discussion and debriefing which lasted about 20 minutes each. The entire simulation session lasted up to 60 minutes.^[20] The fever messages were from and in line with the revised Integrated Management of Childhood Illnesses protocol for Ghana. The video was inspired by Abed *et al.*,^[32] and the scenario and role modeling simulations were designed using a gain-framed message format recommended by Tuong *et al.*^[33] The video could be played on laptops, projected on screens, and on mobile phones. A secure and quiet environment was ensured to facilitate clear viewing, listening, and debriefing. Some participants were educated in groups, while others were taught one-on-one. Interventions such as video, scenario and role modeling was done for mothers only once. After each session, debriefing was carried out to determine and to ensure that each person had completed the task of recalling or knowing information that have been packaged for them and were educated with. Data collection was done before and two weeks after the simulation-based education sessions.

The control group received routine education on fever alongside regular care education from ward staff and data collectors. The education was based on the revised Integrated Management of Childhood Illnesses protocol on febrile illnesses to cover all necessary sections of the questionnaire two weeks after education; the parents were invited for the posttest. There was no lost to follow-up, and no parent received a financial reward.

Data analysis

Data were analyzed using the Statistical Package for Social Sciences (SPSS) version 16, and the Kolmogorov-Smirnov test was used to determine normality of the data. Demographic data of parents and children were analyzed descriptively with means, standard deviations, and frequencies and were compared between the two groups using the Chi-square test. We used the independent

samples *t*-test to compare parents' knowledge, attitude, and management of fever in children between the two groups. The paired *t*-test was also used for within-group comparisons of the parents' mean knowledge, attitude, and management of fever. Analysis of covariance was used to examine the effect of baseline knowledge on the outcomes. A significance level of <0.05 was applied.

Ethical considerations

The study received approval from The Ethics and Institutional Review Board of Tehran University of Medical Sciences (IR.TUMS.FNM.REC.1398.126) and the Department of Research and Development of the Tamale Teaching Hospital (TTH/R&D/SR/117). Parents were briefed about the study objectives and were also provided with written consent for participation in the study. They were also assured about voluntary participation and the confidential management of their personal information.

Results

The mean age of parents in the simulation-based and routine education groups were 29.20 ± 6.45 and 30.13 ± 5.98 years, respectively ($P=0.508$). There were no significant differences in the characteristics of parents and children between the two groups [Table 1].

With the exception of mean baseline knowledge ($P<0.0001$), there were no significant differences in the pretest mean scores between the simulation-based and routine education groups. All the posttest mean scores were better than the pretest in both groups. However, on the posttest, the mean scores for knowledge, attitude, and management were significantly higher in the simulation-based education than in the routine education group ($P<0.05$, [Table 2]).

The study revealed that knowledge scores on the pretest were statistically different between the two groups. However, an analysis of covariance failed to demonstrate its effect on the study outcomes.

Discussion

At the end of the study, parents' knowledge, attitude, and management improved better in the simulation-based education group than in the routine education group. Prior knowledge was not found to affect the study outcomes. These findings imply that, even though both groups experienced improvements, simulation-based education could offer better effects on parents' knowledge, attitude, and management of fever in children.

Table 1. Demographic characteristics of parents and children ^a

Variables	Group		P-value
	Simulation-based education	Routine education	
Parent age, year	29.20±6.45	30.13±5.98	0.508
Age of child, month	32.75±20.35	34.75±26.29	0.705
Child's temperature	38.33±0.52	38.43±0.68	0.431
Marital status			0.147
Married	35(87.5)	27 (67.5)	
Single	3 (7.5)	8 (20.0)	
Divorced	2 (5.0)	3 (7.5)	
Widowed	0 (0.0)	2 (5.0)	
Residency			0.176
Urban	19 (47.5)	26 (65.0)	
Rural	21 (52.5)	14 (35.0)	
Level of education			0.364
Basic level	6 (15.0)	5 (12.5)	
High school	7 (17.5)	14 (35.0)	
Tertiary	9 (22.5)	8 (20.0)	
No education	18 (45.0)	13 (32.5)	
Febrile seizure			0.858
Yes	15(37.5)	13 (32.5)	
No	24 (60.0)	25 (62.5)	
I don't know	1(2.5)	2 (5.0)	

^a Data presented as Mean ±SD or n (%)

Table 2. Comparison of pretest and posttest means scores between simulation-based and routine education groups ^a

Variables	Groups		P-value (t-test)
	Simulation-based education	Routine education	
Knowledge			
Pretest	21.63±2.40	23.98±2.57	<0.0001
Post test	26.38±4.36	24.75±2.72	0.048
Change	4.75±5.41	0.78±2.61	<0.000
P-value (Paired t-test)	<0.000	0.068	
Attitude			
Pretest	28.48±6.03	28.33±5.98	0.911
Post test	39.90±8.33	33.40±3.95	<0.0001
Change	11.43±9.00	5.08±6.90	<0.001
P-value (Paired t-test)	<0.000	<0.000	
Management			
Pretest	25.15±4.23	24.43±3.34	0.400
Posttest	32.33±5.18	30.15±4.12	0.040
Change	7.18±7.14	5.72±4.22	0.272
P-value (Paired t-test)	<0.000	<0.000	

^a Data presented as Mean±SD

The findings of Baker *et al.*,^[26] Prawesti *et al.*,^[34] and Ong *et al.*^[25] support our result. These studies compared the effects of education using videos and brochures on maternal health literacy or knowledge regarding fever. Similar to our study, video education resulted in better

outcomes in terms of maternal health literacy or knowledge about fever control in children. The above studies however recommended both methods as being suitable for educating parents to minimize anxiety and reduce pediatric emergency unit visitation by

parents.^[25,26,34] Video presentations produced for a target population by using the language that parents understand are likely to be easily accessible and comprehensible, especially with the use of social media. Literate parents may also have an added advantage if both methods are combined. This would help parents to make better and more informed decisions related to acute health situations of their children, as reported in some earlier studies.^[13,32,35] In some studies in Iran and Egypt, simulation training was also found to be more effective than routine education in improving parents' knowledge, attitude, and practice regarding the prevention of febrile seizures in children.^[36,37] However, a study of inhaler use skills indicated that both face-to-face and video-based education methods could improve the participants' skills, even though the face-to-face method was more effective.^[38] Despite controversies about the effects of different educational methods, the blended simulation-based method used in the present study seems to combine all the effects of face-to-face, video, role playing, and discussion and helps nurses and other healthcare providers understand parents' worries and problems, alleviate their concerns, focus on their educational needs, receive their feedback, correct their misconceptions, and modify their fever management behavior.

The simulation video was produced by the researcher, without having professional expertise in acting and video editing. This may have affected the implementation and hence the outcome of the study. More standardized videos are needed to guide researchers in the implementation and subsequent evaluation of their effects. Another likely limitation was the sampling technique and sample size. Only 80 subjects were selected and non-randomly assigned to the study groups. This could also be a source of limitation in this study.

Conclusions

Simulation-based education is widely used and very effective depending on the objective and target audience. In our study, we found that visual aids such as videos, discussions in the form of scenarios, and role-playing were effective in educating parents about childhood fever. We also noted that the routine methods could also be useful. Perhaps the routine method would be more effective if it is combined with simulation approaches. We would then recommend the use of methods depending on the design and objective of an education process.

Acknowledgment

The authors wish to thank all the parents who participated in this study.

Competing interests

There are no conflicts of interest.

Abbreviations

Statistical Package for Social Sciences: SPSS.

Authors' contributions

All authors read and approved the final manuscript. All authors take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This study was approved and financially supported by Tehran University of Medical Sciences.

Role of the funding source

None.

Availability of data and materials

The data used in this study are available from the corresponding author on request.

Ethics approval and consent to participate

The study received approval from The Ethics and Institutional Review Board of Tehran University of Medical Sciences (IR.TUMS.FNM.REC.1398.126) and the Department of Research and Development of the Tamale Teaching Hospital (TTH/R&D/SR/117). Parents were briefed about the study objectives and were also provided with written consent for participation in the study.

Consent for publication

By submitting this document, the authors declare their consent for the final accepted version of the manuscript to be considered for publication.

References

1. Niehues T. The febrile child: diagnosis and treatment. *Dtsch Arztebl Int* 2013;110:764-73. doi:10.3238/arztebl.2013.0764 PMID:24290365 PMCID:PMC3849992
2. Roy S, Simalti AK. Comparison of Antipyretic Efficacy of Intravenous (IV) Acetaminophen versus Oral (PO) Acetaminophen in the Management of Fever in Children. *Indian J Pediatr* 2018;85:1-4. doi:10.1007/s12098-017-2457-3 PMID:28887752
3. Chiappini E, Cangelosi AM, Becherucci P, Pierattelli M, Galli L, de Martino M. Knowledge, attitudes and misconceptions of Italian healthcare professionals regarding fever management in children. *BMC Pediatr* 2018;18:1-6. doi:10.1186/s12887-018-1173-0 PMID:29914424 PMCID:PMC6007065

4. Kelly M, Sahn LJ, Shiely F, O'Sullivan R, de Bont EG, Mc Gillicuddy A, et al. Parental knowledge, attitudes and beliefs on fever: a cross-sectional study in Ireland. *BMJ Open* 2017;7:e015684. doi:10.1136/bmjopen-2016-015684 PMID:28694348 PMCID:PMC5541503
5. Thompson AP, Nesari M, Hartling L, Scott SD. Parents' experiences and information needs related to childhood fever: a systematic review. *Patient Educ Couns* 2020;103:750-763. doi:10.1016/j.pec.2019.10.004 PMID:31668490
6. Nyarko SH, Cobblah A. Sociodemographic determinants of malaria among under-five children in Ghana. *Malar Res Treat* 2014;2014:304361. doi:10.1155/2014/304361 PMID:25580349 PMCID:PMC4279724
7. Ojeikere KI. Malaria Prevention and Treatment Seeking Practices among Mothers of Children Under Five Years in Adentan Municipality. University of Ghana; 2016. Available from: <https://ugspace.ug.edu.gh/handle/123456789/21532> [Last access date: 20 Feb 2023]
8. Ansah EK, Gyapong M, Narh-Bana S, Bart-Plange C, Whitty CJ. Factors influencing choice of care-seeking for acute fever comparing private chemical shops with health centres and hospitals in Ghana: a study using case-control methodology. *Malar J* 2016;15:290. doi:10.1186/s12936-016-1351-1 PMID:27225480 PMCID:PMC4880965
9. Nonvignon J, Chinbuah MA, Gyapong M, Abbey M, Awini E, Gyapong JO, et al. Is home management of fevers a cost-effective way of reducing under-five mortality in Africa? The case of a rural Ghanaian District. *Trop Med Int Health* 2012;17:951-957. doi:10.1111/j.1365-3156.2012.03018.x PMID:22643324
10. Baiden F, Owusu-Agyei S, Bawah J, Bruce J, Tivura M, Delmini R, et al. An evaluation of the clinical assessments of under-five febrile children presenting to primary health facilities in rural Ghana. *PLoS one* 2011;6:e28944. doi:10.1371/journal.pone.0028944 PMID:22174932 PMCID:PMC3236777
11. Webster J, Baiden F, Bawah J, Bruce J, Tivura M, Delmini R, et al. Management of febrile children under five years in hospitals and health centres of rural Ghana. *Malar J* 2014;13:261. doi:10.1186/1475-2875-13-261 PMID:25008574 PMCID:PMC4114131
12. Enarson MC, Ali S, Vandermeer B, Wright RB, Klassen TP, Spiers JA. Beliefs and expectations of Canadian parents who bring febrile children for medical care. *Pediatrics* 2012;130:e905-e912. doi:10.1542/peds.2011-2140 PMID:22966028
13. Kelly M, Sahn L, McCarthy S, O'Sullivan R, Mc Gillicuddy A, Shiely F. Randomised controlled trial of an intervention to improve parental knowledge and management practices of fever. *BMC Pediatr* 2019;19:447. doi:10.1186/s12887-019-1808-9 PMID:31739785 PMCID:PMC6863059
14. Barakat K. The role of simulation-based education in cardiology. *Heart* 2019;105:728-732. doi:10.1136/heartjnl-2017-311153 PMID:30661036
15. Baumann SL, Sharoff L, Penalo L. Using Simulation to Enhance Global Nursing. *Nurs Sci Q* 2018;31:374-378. doi:10.1177/0894318418792877 PMID:30223748
16. Brown CE, Back AL, Ford DW, Kross EK, Downey L, Shannon SE, et al. Self-assessment scores improve after simulation-based palliative care communication skill workshops. *Am J Hosp Palliat Care* 2018;35:45-51. doi:10.1177/1049909116681972 PMID:28273752
17. Koivisto JM, Hannula L, Bøje RB, Prescott S, Bland A, Rekola L, et al. Design-based research in designing the model for educating simulation facilitators. *Nurse Educ Pract* 2018;29:206-211. doi:10.1016/j.nepr.2018.02.002 PMID:29433039
18. Crowe S, Ewart L, Derman S. The impact of simulation based education on nursing confidence, knowledge and patient outcomes on general medicine units. *Nurse Educ Pract* 2018;29:70-75. doi:10.1016/j.nepr.2017.11.017 PMID:29190590
19. Shapiro M, Morey J, Small S, Langford V, Kaylor C, Jagminas L, et al. Simulation based teamwork training for emergency department staff: does it improve clinical team performance when added to an existing didactic teamwork curriculum?. *Qual Saf Health Care* 2004;13:417-421. doi:10.1136/qshc.2003.005447 PMID:15576702 PMCID:PMC1743923
20. Chang LC, Lee PI, Guo NW, Huang MC. Effectiveness of simulation-based education on childhood fever management by Taiwanese parents. *Pediatr Neonatol* 2016;57:467-473. doi:10.1016/j.pedneo.2015.10.011 PMID:27036101
21. Chiappini E, Bortone B, Galli L, de Martino M. Guidelines for the symptomatic management of fever in children: systematic review of the literature and quality appraisal with AGREE II. *BMJ open* 2017;7:e015404. doi:10.1136/bmjopen-2016-015404 PMID:28760789 PMCID:PMC5642818
22. Huang MC, Liu CC, Chi YC, Thomas K, Huang CC. Effects of educational intervention on changing parental practices for recurrent febrile convulsions in Taiwan. *Epilepsia* 2002;43:81-86. doi:10.1046/j.1528-1157.2002.35501.x PMID:11879391
23. Kelly L, Morin K, Young D. Improving caretakers' knowledge of fever management in preschool children: is it possible? *J Pediatr Health Care* 1996;10:167-173. doi:10.1016/S0891-5245(96)90040-1 PMID:8920378
24. Wright A, Moss P, Dennis DM, Harrold M, Levy S, Furness AL, et al. The influence of a full-time, immersive simulation-based clinical placement on physiotherapy student confidence during the transition to clinical practice. *Adv Simul (Lond)* 2018;3:3. doi:10.1186/s41077-018-0062-9 PMID:29484204 PMCID:PMC5819286
25. Ong TEL, Kua JPH, Liew LJ, Lim ZY, Thia MXH, Sung SC. Assessing effective methods to educate caregivers on fever in children aimed at reducing input to the paediatric emergency department. *Proc Singapore Healthc* 2018;27:73-84. doi:10.1177/2010105817733271
26. Baker MD, Monroe KW, King WD, Sorrentino A, Glaeser PW. Effectiveness of fever education in a pediatric emergency department. *Pediatr Emerg Care* 2009;25:565-568. doi:10.1097/PEC.0b013e3181b4f64e PMID:19755888
27. Moon RY, Hauck FR, Colson ER, Kellams AL, Geller NL, Heeren T, et al. The effect of nursing quality improvement and mobile health interventions on infant sleep practices: a randomized clinical trial. *JAMA* 2017;318:351-359. doi:10.1001/jama.2017.8982 PMID:28742913 PMCID:PMC5593130
28. Service GS. Ghana demographic and health survey 2014. In.: GSS, GHS, and ICF International Rockville, Maryland, USA; 2015.
29. Chiappini E, Parretti A, Becherucci P, Pierattelli M, Bonsignori F, Galli L, et al. Parental and medical knowledge and management of fever in Italian pre-school children. *BMC Pediatr* 2012;12:97. doi:10.1186/1471-2431-12-97 PMID:22794080

PMCID:PMC3439692

30. Chiappini E, Cangelosi AM, Becherucci P, Pierattelli M, Galli L, de Martino M. Knowledge, attitudes and misconceptions of Italian healthcare professionals regarding fever management in children. *BMC Pediatr* 2018;18:194. doi:10.1186/s12887-018-1173-0 PMid:29914424 PMCID:PMC6007065
31. Greensmith L. Nurses' knowledge of and attitudes towards fever and fever management in one Irish children's hospital. *J Child Health Care* 2013;17:305-316. doi:10.1177/1367493512461457 PMid:23455871
32. Abu Abed M, Himmel W, Vormfelde S, Koschack J. Video-assisted patient education to modify behavior: A systematic review. *Patient Educ Couns* 2014;97:16-22. doi:10.1016/j.pec.2014.06.015 PMid:25043785
33. Tuong W, Larsen ER, Armstrong AW. Videos to influence: a systematic review of effectiveness of video-based education in modifying health behaviors. *J Behav Med* 2014;37:218-233. doi:10.1007/s10865-012-9480-7 PMid:23188480
34. Prawesti I, Haryanti F, Lusmilasari L. Effect of health education using video and brochure on maternal health literacy. *Belitung Nurs J* 2018;4:612-618. doi:10.33546/bnj.176
35. Cook DA, Andersen DK, Combes JR, Feldman DL, Sachdeva AK. The value proposition of simulation-based education. *Surgery* 2018;163:944-949. doi:10.1016/j.surg.2017.11.008 PMid:29452702
36. Najimi A, Dolatabadi NK, Esmaeili AA, Sharifirad GR. The effect of educational program on knowledge, attitude and practice of mothers regarding prevention of febrile seizure in children. *J Educ Health Promot* 2013;2:26. doi:10.4103/2277-9531.112703 PMid:24083276 PMCID:PMC3778569
37. Tavan A, Monemi E, Keshavarz F, Kazemi B, Nematollahi M. The effect of simulation-based education on parental management of fever in children: a quasi-experimental study. *BMC Nurs* 2022;21:168. doi:10.1186/s12912-022-00938-7 PMid:35764950 PMCID:PMC9238067
38. Adib-Hajbaghery M, Karimi Z. Comparing the effects of face-to-face and video-based education on inhaler use: A randomized, two-group pretest/posttest study. *Iran J Nurs Midwifery Res* 2018;23:352-357 doi:10.4103/ijnmr.IJNMR_17_18 PMid:30186339 PMCID:PMC6111659

How to Cite this Article:

Pouraboli B, Safaiee Fakhr A, Begjani J, Hrdi AM, Ranjbar H, Heidari ME. Effects of simulation-based education on Ghanaian parents' knowledge, attitude and management of fever control. *Nurs Midwifery Stud* 2023;12(3):135-141. doi:10.48307/NMS.2023.407976.1233