Research Article



The relationship between resilience, quality of life, and body image in women with breast cancer undergoing chemotherapy

Faezehsadat Mousavi 10¹, Masoud Bahrami 10¹, Elaheh Ashouri 10^{*1}

¹Nursing and Midwifery Care Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

* **Corresponding author:** Elaheh Ashouri, Nursing and Midwifery Care Research Center, Isfahan University of Medical Sciences, Isfahan, Iran. **Email:** ashori@nm.mui.ac.ir

Received: 21 June 2023 Revised: 21 July 2023 Accepted: 28 July 2023 e-Published: 22 September 2023

Abstract

Background: The diagnosis and treatment of breast cancer put women at risk of altered body image and quality of life (QOL). **Objectives:** This study aimed to investigate the relationship between resilience, QOL, and body image in women with breast cancer undergoing chemotherapy.

Methods: This cross-sectional study was conducted in 2022 on 172 women with breast cancer referring to the chemotherapy department of Omid Hospital in Isfahan, Iran. Data were collected using a demographic data form, the Connor-Davidson Resilience Scale, the World Health Organization Quality of Life questionnaire (WHOQOL-BREF), and the Hopwood Body Image Scale (HBIS). The data were analysed using one-way analysis of variance, Pearson's correlation, Spearman's correlation, and multiple linear regression analysis.

Results: The mean total QOL score was 53.70 ± 18.94 . In addition, the mean resilience score was 69.83 ± 14.52 . The mean body image score was 8.23 ± 6.65 . Resilience was directly correlated with QOL (r=0.491, P<0.001). The psychological-spiritual dimension of QOL was a significant predictor for resilience (P<0.001). Resilience was inversely correlated with body image (r=-0.37, P<0.05).

Conclusion: The psychological-spiritual dimension of QOL was a significant predictor for resilience. Improving women's resilience during chemotherapy can reduce their psychological symptoms and improve the QOL and body image of these patients.

Keywords: Breast cancer, Chemotherapy, Psychological Resilience, Quality of life, Body image.

Introduction

Breast cancer (BC) is the most common cancer in women,^[1] with a prevalence of 11.7% among all cancers.^[2] The number of people with BC is predicted to increase from 2.26 million in 2020 to more than 3.19 million in 2040.^[3] BC is also one of the most prevalent malignancies in Iran,^[4] accounting for 28.1% of all cancers in Iranian women.^[5] Women with BC receive various treatments, including surgery, chemotherapy, and radiotherapy.^[6] Chemotherapy is a systemic method that affects not only malignant cells, but also other rapidly proliferating cells in the body, such as hair follicles, bone marrow, and the digestive and reproductive systems.^[7] In addition, chemotherapy negatively affects the patient's self-concept, body image (BI), sexual relationships, and QOL. The cancer diagnosis, along with the significant side effects of chemotherapy, leads to physical, functional, psychological, and social problems and disrupts the patient's QOL.^[8]

QOL is an individual's perception of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectations, standards, and concerns.^[9] Since the number of BC survivors is increasing, QOL is a major concern for these patients.^[10] Furthermore, women with BC experience a distorted BI.

BI is a set of conscious and unconscious attitudes that people have toward their bodies and is a main factor affecting QOL.[11] A recent study reported that women who had undergone mastectomy experienced negative BI, QOL, and emotional poor distress during chemotherapy.^[12] Another study also reported that chemotherapy lowered the QOL of women with BC due to changes in their BI.^[13] A study also confirmed the relationship between BI and QOL in patients with BC, and identified body dissatisfaction as the strongest predictor of QOL.^[14] Therefore, identifying factors affecting patients'

Copyright© 2023. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms. Downloaded from: https://nmsjournal.kaums.ac.ir/

Mousavi et al

QOL and BI may provide treatment teams with new solutions to help patients with BC.

Resilience protects a person while recovering from difficult conditions such as cancer.^[15] Resilience is a dynamic process that encourages a person to successfully cope with complex or challenging life experiences^[16] Resilience is of particular importance in severe diseases such as cancer. It is believed that an individual's resilience can be strengthened. Improved resilience is associated with improved health, alleviation of illness, and accelerated recovery.^[15] A study found that women with newly diagnosed breast cancer were less resilient, and that resilience was directly correlated with QOL.^[17] However, another study found no significant correlation between resilience and QOL in patients with BC.^[18]

Given the impaired QOL and BI in patients with BC and the controversy over the association between resilience, QOL, and BI, the question arises whether resilience, QOL, and BI are correlated in women with BC undergoing chemotherapy.

Objectives

This study aimed to determine the relationship between resilience, QOL, and BI in women with BC undergoing chemotherapy.

Methods

Study design and participants

A cross-sectional study was conducted between November 2021 and March 2022 on 172 women with BC admitted to the chemotherapy department of Omid Hospital in Isfahan, Iran. The sample size was calculated based on the results of a former study.^[19] Then, with a type I error of 0.05, Z1=1.96, Z2=0.8, r=0.22, and using the following formula $[n = \frac{(z_1+z_2)^2(1-r^2)}{r^2} + 2]$, with a 10% chance of falling, the sample size was calculated at 172. Convenience sampling was used to recruit eligible subjects. Eligible women were recruited by the researcher (F.S, M) based on the file review.

After obtaining written informed consent, the researcher provided the participants with questionnaires to personally complete in a private setting. If the patients were unable to complete the questionnaire due to illiteracy or their illness, the researcher read the questionnaire items to them and recorded their answers on the questionnaire.

Inclusion criteria included a definite medical diagnosis of BC, undergoing breast surgery, undergoing inpatient or outpatient chemotherapy, passing at least three months since the diagnosis of BC, no history of disease recurrence (being in the first phase of diagnosis and treatment of the disease), and willingness to participate in the study. The only exclusion criterion was an incomplete response to the questionnaire.

Data collection instruments

We used a four-part instrument to collect the study data. The instruments included a demographic data form, the Connor-Davidson Resilience Scale (CD-RISC), the World Health Organization quality of Life questionnaire (WHOQOL-BREF), and the Hopwood Body Image Scale (HBIS). The demographic data form included questions on the participant's age, marital status, number of children, education level, occupation, spouse's education, spouse's occupation, place of residence, time passed from the disease diagnosis, chemotherapy phase, previous treatments, and type of surgery.

The WHOQOL-BREF questionnaire contains 26 items in 4 subscales namely, physical health, psychological health, social relationships, and environmental health. All items are answered on a five-point Likert scale ranging from 1 to 5 (except for items 3, 4, and 26 that are scored reversely). Scores of each subscale are converted to a standard score ranging from 0 to 100. Higher scores indicate better QOL.^[20] Nejat *et al.* have tested the reliability and validity of the Persian version of WHOQOL-BREF, and the reliability coefficient was reported to be 0.77, 0.77, 0.75, and 0.84 for different subscales.^[21]

The HBIS contains 10 items arranged on a 4-point Likert scale (not at all=0 to very much=3) and briefly evaluates emotional issues (feeling self-conscious), behavioral issues (difficulty looking at the naked body) and cognitive issues (such as self-satisfaction). This instrument was designed to measure the BI of cancer patients. The minimum and maximum scores are 0 and 30, respectively, with higher scores indicating more symptoms and higher distress or concern about BI.^[22] Rajabi *et al.* reported a reliability coefficient of 0.70 for this scale.^[23]

The CD-RISC contains 25 items scored on a 5-point Likert scale ranging from zero (not true at all) to four (true nearly all of the time). The total score ranges from 0 to 100, with higher scores indicating greater resilience.^[24] Amini *et al.* reported an internal consistency coefficient of 0.83 for this scale.^[25]

Data analysis

Data analysis was performed using the SPSS software version 16. Descriptive statistics (frequency, percent, mean, and standard deviation) were used to describe participants' characteristics. Pearson's or Spearman's

Resilience, Quality of Life and Body Image in Women with Breast Cancer

correlation coefficients were used to examine the correlation between the main variables.

The independent samples t-test, and one-way analysis of variance were used to compare mean BI, QOL, and resilience scores between subgroups of participants.

Multiple linear regression analysis was performed to predict the impact of QOL and BI subscale scores on resilience. The significance level was set at P<0.05.

Ethical considerations

This study was approved by the Ethics Committee of Isfahan University of Medical Sciences under the ethics code IR.MUI.NUREMA.REC.1400.147. Before the questionnaires were distributed, the researcher explained the study objectives and obtained informed consent from the patients. All participants were assured of data confidentiality and were free to participate or withdraw from the study.

Results

The mean age of the patients was 49.65 ± 11.11 years. The mean chemotherapy phase was 9.25 ± 5.75 , and on average, 8.24 ± 3.85 months had passed since the diagnosis of BC. Other demographic information is presented in [Table 1]. The total mean QOL score of women with BC was 53.70 ± 18.94 , with the highest and the lowest mean scores were related to the environmental (56.56 ± 13.83) and the physical (47.98 ± 19.07) subscales, respectively. The mean resilience score was 69.83 ± 14.52 , and the mean BI score was 8.23 ± 6.65 [Table 2].

	Table 1. The demographic characterist	ics of the participants and the	ir mean quality of life, body image	, and resilience scores
--	---------------------------------------	---------------------------------	-------------------------------------	-------------------------

Variable		n (%)	Quality of life	Body image	Resilience
Marital status					
	Single	9 (5.2)	63.88±13.17	5.88 ± 6.60	71.75 ± 21.05
	Married	144 (83.7)	60.50±19.59	8.69±6.56	69.72±14.39
	Divorced	3 (1.9)	58.33 ± 14.43	16.66±6.65	61.80±8.09
	Widowed	16 (9.2)	64.06 ± 17.00	3.76 ± 4.75	71.26±12.89
Number of					
children	Without children	18 (10.5)	63.19±13.19	7.44 ± 6.94	68.41±17.90
	1 child	20 (11.6)	56.25 ± 23.47	9.84±8.95	70.16 ± 14.72
	2 children	62 (36)	63.70±21.91	8.80 ± 6.24	68.46±14.15
	More than 2 children	72 (41.9)	59.37±15.66	7.47±6.17	71.26 ± 14.01
Education					
	Illiterate	58 (33.7)	58.72 ± 18.40	9.18±6.97	69.43±14.34
	Sub-diploma and diploma	87 (50.6)	61.02±19.13	8.41±6.39	68.83±14.67
	Associate or Bachelor's degrees	22 (12.8)	59.09 ± 26.86	8.62±6.66	71.49 ± 11.04
	Above bachelor's degree	5 (2.9)	68.75±8.83	4.50 ± 6.36	88.54±2.94
Occupation					
	Employed	12 (7)	60.78 ± 4.86	6.08 ± 4.96	79.34±10.40
	Housekeeper	150 (87.2)	52.97±12.71	8.43 ± 6.74	69.33±14.60
	Retired	10 (5.8)	56.13±9.89	7.70 ± 7.06	65.90±14.26
Place of residence					
	Urban area	87 (50.6)	55.91±11.02	8.22±6.66	70.85±13.96
	Rural area	85 (49.4)	51.43 ± 13.20	8.23±6.67	68.79±15.08
Spouse's education					
	Illiterate	50 (29.1)	58.72 ± 18.40	9.18±6.97	69.43±14.34
	Sub-diploma and diploma	107 (62.2)	61.02±19.13	8.41±6.39	68.83±14.67
	Associate or and Bachelor's degrees	13 (7.5)	59.09 ± 26.86	8.62±6.66	71.49 ± 11.04
	Above bachelor's degree	2 (1.2)	68.75±8.83	4.50 ± 6.36	88.54±2.94
Spouse's					
occupation	Unemployed	11 (6.4)	63.75±14.96	7.80 ± 5.37	68.08 ± 21.50
	Employee	26 (15.1)	59.65±19.64	7.71±4.45	65.53±16.65
	Self-employed	72 (41.9)	61.08 ± 21.83	9.24±6.86	67.78±13.89
	Teacher	6 (3.5)	65.00 ± 10.45	4.80 ± 5.06	72.70 ± 17.51
	Other	57 (33.1)	58.67±17.89	8.91±7.19	73.46±11.26

Nurs Midwifery Stud. 2023;12(3):151-157 | 153

Mousavi et al

Pearson's correlation coefficient showed that the resilience score was directly associated with the total QOL score (r=0.491, P<0.001) and all its subscales, with the strongest correlation observed for the psychological-spiritual subscale. An inverse correlation was observed between the BI scores and all QOL scores [Table 3]. Furthermore, an inverse correlation was found between resilience and overall BI score (r=-0.37, P<0.05).

Table 2. The mean scores for quality of life and its subscales, resilience, and body image

Variables	Mean±SD
Quality of life subscales	
Physical	47.98±19.07
psychological-spiritual	54.91±15.81
Social relationships	53.39±23.54
Environment	56.56±13.83
Total score	53.70±12.32
Resilience	69.83±14.52
Body image	8.23±6.65

Multiple linear regression analysis showed that among the QOL subscales and the overall BI score, only the psychological-spiritual subscale score significantly predicted resilience [Table 4].

In addition, Pearson's correlation coefficient showed that age and duration of chemotherapy phase were not significantly correlated with resilience, body image, and total QOL scores (P>0.05). The time passed since being informed of the disease correlated directly with the BI score and inversely with the total QOL score (P<0.05). The total QOL score was also directly associated with education level of patients and their spouses (P<0.05) [Table 5].

Furthermore, the mean BI score was significantly higher in married women than in unmarried women (P<0.05). The mean resilience score was significantly higher in employed women than in housekeeper and retired women (P<0.05). Besides, the mean QOL score was significantly higher in women who lived in urban areas than in women who lived in rural areas (P<0.05) [Table 6].

Table 3. Pearson correlation coefficients of resilience scores and body image with the total score of quality of life and its subscales

Quality of life subscales	Resilience score		Body i	mage score
	r	Р	r	Р
Physical	0.194	0.01	-0.279	< 0.001
Psychological-spiritual	0.637	< 0.001	-0.491	< 0.001
Social Relationships	0.241	0.001	-0.237	0.002
Environment	0.339	< 0.001	-0.222	0.003
Total Score	0.491	< 0.001	-0.415	< 0.001

Table 4. Results of regression analysis for predicting the impact of dimensions of quality of life and body image on resilience

Score	Raw coefficient	Standardized coefficient	Т	P-value
Physical	-0.071	-0.094	1.39	0.17
Psychological-spiritual	0.551	0.600	7.60	< 0.001
Social relationships	-0.002	-0.003	0.05	0.96
Environment	0.070	0.067	0.94	0.35
Body image	-0.190	-0.087	1.27	0.20

 Table 5. Correlation between some demographic parameters, resilience, body image, and total quality of life score

Variable	Resilience Score		Body Ima	Body Image Score		Total quality of life scores	
	r	Р	r	Р	r	Р	
Age	-0.065	0.39	-0.032	0.68	-0.136	0.08	
Chemotherapy phase	0.064	0.40	0.065	0.39	-0.148	0.06	
The duration of being	-0.124	0.10	0.163	0.03	-0.229	0.003	
informed about the disease							
Number of children	0.074	0.34	-0.051	0.51	-0.014	0.85	
Patients education level	0.079	0.30	-0.120	0.12	0.220	0.004	
Spouse's education level	0.036	0.66	-0.052	0.53	0.159	0.04	

Resilience, Quality of Life and Body Image in Women with Breast Cancer

Table 6. The independent samples <i>t</i> -test of Urban and Rural women with resilience, body image quality of life							
Score	Urban Rural The independent samples						
-	Mean ± SD	Mean ± SD	t	df	Р		
Resilience	70.85±13.96	68.79±15.08	0.93	170	0.35		
Body Image	8.22±6.66	8.23±6.67	0.008	170	0.99		
Quality of life	55.91±11.02	51.43 ± 13.20	2.42	170	0.02		

Discussion

The women who participated in the present study scored about 50% of the possible QOL score. They also possessed the highest and lowest QOL scores in the environmental and the physical subscales, respectively. These findings are consistent with some of the previous studies.^[26,27]

The mean resilience score in the present study was higher than those of women with BC in two earlier studies conducted in Tehran, Iran ^[28] and in China.^[17] However, in a previous study conducted in Omid Hospital in Isfahan, women with BC achieved higher resilience scores than in our study.^[29]

This difference could be due to the complexity of this concept, since several parameters such as spirituality and culture-related activities, may affect individual's resilience.

In the current study, women with BC scored lower BI than participants in a recent study conducted in Tehran, Iran.^[30] This difference may be attributed to the duration of the disease. The present study was conducted in women with newly diagnosed BC who had recently undergone surgery and chemotherapy. However, the latter study was conducted with women who had been diagnosed with BC for at least one year and who had become somewhat accustomed to changes in their BI.

In the present study, resilience scores of women with BC were directly correlated with total scores as well as the scores of all QOL subscales. Moreover, linear regression analysis showed that of the QOL subscales and the total BI score; only the psychological-spiritual subscale scores could predict resilience. These findings are consistent with what was reported by Tu et al. who studied the role of trait resilience and coping styles in positive psychological changes after the diagnosis and treatment of BC.^[31] Zhou et al. also reported that resilience indirectly affects QOL through social support and positive coping styles.^[32] High resilience plays a vital role in positive psychological changes and improving the psychological dimensions of QOL.^[31] However, a study by Boškailo et al, found no significant correlation between resilience and QOL in women with BC.^[18] The insignificant correlation in the latter study might be attributable to the fact that it was conducted with only 60 patients.

However, our study had a larger sample size, so its results

may be more reliable.

The findings of the present study showed an inverse association between resilience and BI scores. Our findings are consistent with the results of Hsu et al. who studied the relationship between BI and resilience in patients with BC.^[33] Patients with low resilience appear to have a more pessimistic attitude toward their bodies, and resilience is a crucial protective factor for maintaining a positive BI. In this study, resilience was not significantly associated with age, education level, and spouse's education level. However, employed women exhibited significantly higher levels of resilience than housekeeper and retired women. Plitzko et al. also reported that unemployed patients with poor socioeconomic status had lower levels of resilience.^[34] However, another study in the United States found no relationship between employment and resilience among women.^[35]

These contradictory results indicate that resilience is a personal trait that might be affected by several factors such as culture.

In the current study, the total QOL score was directly correlated with the education levels of patients and their husbands. Considering the role of socioeconomic and cultural factors in health issues and the link between QOL and health, it is possible to justify a higher QOL for people with a higher education level.

Our findings showed that time passed since diagnosis of BC was directly correlated with BI, whereas it was indirectly associated with QOL. These findings are consistent with those of Wu *et al.* who studied changes of BI and QOL in patients with BC.^[36] QOL appears to be affected by the duration of illness. Thus, women have a higher QOL in the early stages of disease diagnosis, but the QOL declines over time as the disease and treatment complications emerge.

In this study, mean QOL scores were significantly higher for women living in urban areas than for women living in rural areas. Urban women have more opportunities to participate in cultural and social activities, which appear to have a positive impact on their QOL.

This study had some limitations. First, because we used self-report instruments, the patient's physical and mental state may have influenced the accuracy of their responses.

Mousavi et al

Second, the participants were women with newly diagnosed BC who had recently undergone chemotherapy, and this may limit the generalizability of the results.

Conclusions

This study found a direct correlation between resilience and overall QOL and all its dimensions, and an inverse correlation with BI. The psychological-spiritual dimension of QOL was a significant predictor for resilience. Therefore, nurses and other health care providers are recommended to develop programs that strengthen the psychological resilience in women with BC to reduce their psychological symptoms, and improve their QOL and BI.

Acknowledgment

This article was derived from a master's thesis in Nursing with project number 3400600, Isfahan University of Medical Sciences, Isfahan, Iran. The authors would like to acknowledge the research deputy at Isfahan University of Medical Sciences for their support. We also are thankful for all patients who participated in this study.

Competing interests

There are no conflicts of interest.

Abbreviations

body image: BI Breast cancer: BC quality of life: QOL Hopwood Body Image Scale: HBIS

Authors' contributions

All authors read and approved the final manuscript. All authors take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This study was financially supported by the Isfahan University of Medical Science.

Role of the funding source

None.

Availability of data and materials

The data used in this study are available from the corresponding author on request.

Ethics approval and consent to participate

This study was approved by the Ethics Committee of Isfahan University of Medical Sciences under the ethics code IR.MUI.NUREMA.REC.1400.147. Before the questionnaires were distributed, the researcher explained the study objectives and obtained informed consent from the patients. All participants were assured of data confidentiality and were free to participate or withdraw from the study.

Consent for publication

By submitting this document, the authors declare their consent for the final accepted version of the manuscript to be considered for publication.

References

- Gradishar WJ, Anderson BO, Abraham J, Aft R, Agnese D, Allison KH, *et al.* Breast Cancer, Version 3.2020, NCCN Clinical Practice Guidelines in Oncology. J Natl Compr Canc Netw 2020;18:452-78. doi:10.6004/jnccn.2020.0016 PMid:32259783
- 2. Globocan. Cancer Today. Global Cancer Observatory. Retrieved July 09, 2020, Available from: https://gco.iarc.fr/today/data/factsheets/cancers/20-Breast-factsheet.pdf [Last access date: December, 30 2022]
- 3. Globocan. Cancer Tomorrow. Global Cancer Observatory. Retrieved July 09, 2020, Available from:https://gco.iarc.fr/tomorrow/en/dataviz/isotype?cancers=20 &single_unit=100000 [Last access date: December 30 2022]
- Alizadeh M, Ghojazadeh M, Piri R, Mirza-Aghazadeh-Attari M, Mohammadi S, Naghavi-Behzad M. Age at Diagnosis of Breast Cancer in Iran: A Systematic Review and Meta-Analysis. Iran J Public Health 2021;50:1564-76. doi: 10.18502/ijph.v50i8.6802PMid:34917527 PMCid:PMC8643522
- Globocan. Iran, Islamic Republic of Fact Sheets. Global Cancer Observatory. Retrieved July 10, 2020, Available from:https://gco.iarc.fr/today/data/factsheets/populations/364iran-islamic-republic-of-fact-sheets.pdf [Last access date: December, 30 2022]
- Matzo M, SHerman DW. Palliative Care Nursing: Quality Care to the End of Life. 5th ed: Springer Publishing Company; 2019. doi: 10.1891/9780826127198
- 7. Wilkes GM, Barton-Burke M. 2020-2021 Oncology Nursing Drug Handbook. 23rd ed: Jones & Bartlett Learning; 2020.
- Hinkle JL, H.Cheever K, Overbaugh K. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed: LWW; 2022.
- The World Health Organization Quality of Life assessment (WHOQOL): position paper from the World Health Organization. Soc Sci Med 1995;41:1403-9. doi:10.1016/0277-9536(95)00112-K PMid:8560308
- van Leeuwen M, Husson O, Alberti P, Arraras JI, Chinot OL, Costantini A, et al. Understanding the quality of life (QOL) issues in survivors of cancer: towards the development of an EORTC QOL cancer survivorship questionnaire. Health Qual Life Outcomes 2018;16:114. doi:10.1186/s12955-018-0920-0 PMid:29866185 PMCid:PMC5987570
- 11. Berman A, Snyder S, Frandsen G. Kozier & Erb's Fundamentals of Nursing Concepts, Process, and Practice. 11th ed: Pearson 2022.
- Pereira M, Moreira CS, Nogueira-Silva C, Izdebski P, Pereira MG. Breast cancer post-surgical impact on women's quality of life during chemotherapy treatment: A structural equation modelling approach. Eur J Cancer Care (Engl) 2021;30:e13349. doi:10.1111/ecc.13349
- 13. Phoosuwan N, Lundberg PC. Psychological distress and healthrelated quality of life among women with breast cancer: a

 descriptive
 cross-sectional
 study.
 Support
 Care
 Cancer

 2022;30:3177-86.
 doi:10.1007/s00520-021-06763-z
 PMid:34950961
 PMCid:PMC8857009

- 14. Ettridge K, Scharling-Gamba K, Miller C, Roder D, Prichard I. Body image and quality of life in women with breast cancer: Appreciating the body and its functionality. Body Image 2022;40:92-102. pMid:34902783
- Babić R, Babić M, Rastović P, Ćurlin M, Šimić J, Mandić K, *et al.* Resilience in Health and Illness. Psychiatr Danub 2020;32:226-32.
- 16. Sisto A, Vicinanza F, Campanozzi LL, Ricci G, Tartaglini D, Tambone V. Towards a Transversal Definition of Psychological Resilience: A Literature Review. Medicina (Kaunas) 2019;55:745. doi:10.3390/medicina55110745 PMid:31744109 PMCid:PMC6915594
- Zhou K, Ning F, Wang X, Wang W, Han D, Li X. Perceived social support and coping style as mediators between resilience and health-related quality of life in women newly diagnosed with breast cancer: a cross-sectional study. BMC Womens Health 2022; 22:198. doi:10.1186/s12905-022-01783-1 PMid:35643446 PMCid:PMC9148478
- Boškailo E, Franjić D, Jurić I, Kiseljaković E, Marijanović I, Babić D. Resilience and Quality of Life of Patients with Breast Cancer. Psychiatr Danub 2021;33:572-9.
- 19. Ristevska-Dimitrovska G, Filov I, Rajchanovska D, Stefanovski P, Dejanova B. Resilience and Quality of Life in Breast Cancer Patients. Open Access Maced J Med Sci 2015;3:727-31. doi:10.3889/oamjms.2015.128 PMid:27275317 PMCid:PMC4877917
- 20. Development of the World Health Organization WHOQOL-BREF Quality of Life Assessment. Psychol Med 1998;28:8-551. doi:10.1017/S0033291798006667 PMid:9626712
- 21. Nejat S, Montazeri A, Holakouie Naieni K, Mohammad K, Majdzadeh SR. The World Health Organization quality of Life (WHOQOL-BREF) questionnaire: Translation and validation study of the Iranian version. J School of Public Health and Institute Public Health Res 2006;4:1-12.
- Hopwood P, Fletcher I, Lee A, Al Ghazal S. A body image scale for use with cancer patients. Eur J Cancer 2001;37:189-97. doi:10.1016/S0959-8049(00)00353-1 PMid:11166145
- 23. Rajabi G, Kaveh Farsani Z, Fadaei Dehcheshmeh H, Jelodari A. Psychometric Properties of the Persian version Scale of Body Image among Patients with Breast Cancer. Iran Breast Dis Q 2015; 8:66-74.
- 24. Connor KM, Davidson JR. Development of a new resilience scale: the Connor-Davidson Resilience Scale (CD-RISC). Depress Anxiety 2003;18:76-82. doi:10.1002/da.10113 PMid:12964174
- 25. Amini F. The relationship between resilience and burnout of nurses. J Res Dev Nurs Midw 2013;10:94-102.
- 26. Schröder ML, Stöckigt B, Binting S, Tissen-Diabaté T, Bangemann N, Goerling U, *et al.* Feasibility and Possible Effects of Mindful Walking and Moderate Walking in Breast Cancer Survivors: A Randomized Controlled Pilot Study With a Nested Qualitative Study Part. Integr Cancer Ther 2022;21:15347354211066067. doi:10.1177/15347354211066067 PMid:35045736 PMCid:PMC8777370
- 27. Gangane N, Khairkar P, Hurtig AK, San Sebastián M. Quality of Life Determinants in Breast Cancer Patients in Central Rural

India. Asian Pac J Cancer Prev 2017;18:3325-32.

- Farahbakhshbeh S, Nejad SAM, Moazedian A. Predicting Self-Efficacy of Women with Breast Cancer based on Quality of Life, Religious Orientation, Resilience, Death Anxiety, Psychological Hardiness and Perceived Social Support. Iran J Health Psychol 2019;2:65-78.
- Taheri A, Falavarjani MF. Hope, self-efficacy, and perceived stress among Iranian women with breast cancer: Examining the mediating role of resilience. Middle East J Posit Psychol 2019;5:82-96
- 30. Kiaei T, Makvandi B, Manesh FM, Hafezi F. The causal relationship between perceived stress, perceived social support, and resilience with emotional adaptation mediated by bodyimage of breast cancer patients. Razavi Int J Med 2021;9:81-6.
- Tu PC, Yeh DC, Hsieh HC. Positive psychological changes after breast cancer diagnosis and treatment: The role of trait resilience and coping styles. J Psychosoc Oncol 2020;38:156-70. doi:10.1080/07347332.2019.1649337 PMid:31625826
- 32. Zhou K, Ning F, Wang W, Li X. The mediator role of resilience between psychological predictors and health-related quality of life in breast cancer survivors: a cross-sectional study. BMC Cancer 2022;22:57. doi:10.1186/s12885-022-09177-0 PMid:35022019 PMCid:PMC8756657
- Hsu H-T, Juan C-H, Chen J-L, Hsieh H-F. Mediator Roles of Social Support and Hope in the Relationship Between Body Image Distress and Resilience in Breast Cancer Patients Undergoing Treatment: A Modeling Analysis. Front Psychol 2021;12:695682. doi:10.3389/fpsyg.2021.695682 PMid:34630208 PMCid:PMC8498031
- 34. Plitzko L, Mehnert-Theuerkauf A, Götze H. Resilience in Long-Term Cancer Survivors - Associations with Psychological Distress and Sociodemographic Characteristics. Psychother Psychosom Med Psychol 2020;70:182-9. doi:10.1055/a-0927-6782 PMid:31315142
- Bazzi AR, Clark MA, Winter MR, Ozonoff A, Boehmer U. Resilience Among Breast Cancer Survivors of Different Sexual Orientations. LGBT Health 2018;5:295-302. doi:10.1089/lgbt.2018.0019 PMid:29878863 PMCid:PMC6913098
- Wu TY, Chang TW, Chang SM, Lin YY, Wang J, Kuo YL. Dynamic Changes of Body Image and Quality of Life in Breast Cancer Patients. Cancer Manag Res 2019;11:10563-10571 doi:10.2147/CMAR.S223314 PMid:31908528 PMCid:PMC6925559

How to Cite this Article:

Mousavi F, Bahrami m, Ashouri E. The Relationship between resilience, quality of life, and body image in women with breast cancer undergoing chemotherapy. Nurs Midwifery Stud. 2023;12(3):151-157. doi:10.48307/NMS.2023.179816