



The relationship between functional independence and life satisfaction among Iranian community-dwelling older adults during the COVID-19 pandemic: A cross-sectional study

Mohammad Hossein Saghaei ¹, Mohammad Saleh Poor Emran ², Nima Sarouei ¹, Morteza Zarekar ¹, Shima Haghani ³, Shahzad Pashaeypoor ^{4*}

¹ Student Research Committee, Tehran University of Medical Sciences, Tehran, Iran

² Department of Community Health Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

³ Nursing Care Research Center, Iran University of Medical Sciences, Tehran, Iran

⁴ Department of Community Health and Geriatric Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

* **Corresponding author:** Shahzad Pashaeypoor, Department of Community Health and Geriatric Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran. **Email:** sh-pashaeypour@tums.ac.ir

Received: 4 April 2023 **Revised:** 28 June 2023 **Accepted:** 1 July 2023 **e-Published:** 22 September 2023

Abstract

Background: The COVID-19 pandemic has fundamentally changed the lives of older adults. However, the functional independence (FI) and the life satisfaction (LS) of older adults have received less attention.

Objectives: This study aimed to evaluate the relationship between FI and LS in community-dwelling older adults during the COVID-19 pandemic.

Methods: This cross-sectional study was conducted in 2021. Participants were 479 community-dwelling older adults randomly selected from public healthcare centers in southern Tehran, Iran. Data collection instruments were a demographic questionnaire, the activities of daily living (ADL) scale, the Instrumental ADL (IADL) scale, and the Life Satisfaction Index-Z. The Pearson correlation coefficient, independent samples t-test, analysis of variance, and regression analysis were used to analyze the data.

Results: The mean scores of participants' LS, FI in ADL, and FI in IADL were 11.94 ± 5.11 (in the possible range of 0–26), 15 ± 2.32 (in the possible range of 0–16), and 11.28 ± 3.72 (in the possible range of 0–14), respectively. Around 55% of participants had low LS and only 2.5% of them had high LS. LS had a significant correlation with independence in ADL ($r=0.133$; $P=0.004$) and IADL ($r=0.213$; $P<0.001$). Independence in IADL and income significantly predicted 30% of the variance of LS ($P<0.05$).

Conclusion: Older adults had optimal levels of FI in ADL and IADL and a low level of LS. Healthcare authorities can improve older adults' LS through strategies to promote their FI; such as formulating health policies, amending health system laws, and establishing insurance coverage to reduce costs.

Keywords: Aging, Functional independence, Life satisfaction.

Introduction

Aging is a natural process with extensive physiological, psychological, and social changes.^[1] The population over the age of 65 is expected to double in the next forty years, particularly in developing countries.^[2] The aging population in Iran is estimated to increase from 14% in 2011 to 34% by 2050.^[3]

Physical and mental disabilities, and chronic diseases make older adults unable to perform daily activities,

undermine their authority, reduce their capacity for self-care and self-efficacy, and endanger their functional independence (FI).^[4,5]

FI is the ability to perform activities of daily living (ADL) and instrumental ADL (IADL). ADL refers to self-care activities that should be performed during in living, including eating, clothing, bathing, moving, elimination, and walking. IADL also includes activities needed for independent living in society, such as food preparation,

household and outdoor activities, medication taking, payment management, and telephone use.^[4] The ability to perform ADL and IADL and maintain FI is so important for older adults that they consider dependence as worse than death.^[6] The prevalence of dependence in ADL and IADL among Iranian older adults was reported to be 23.3% and 28.5%, respectively.^[7] A study in Iran also showed that the occurrence of traumatic events in older adults has a significant impact on their ADL.^[8]

One of the most important factors affecting older adults is infectious diseases, such as the current coronavirus disease 2019 (COVID-19) pandemic. Epidemics can impair normal functioning, cause physical weakness, and compromise mental and social health due to quarantine and physical distancing.^[9,10] Moreover, affliction by COVID-19 has significant effects on older adults functioning.^[11,12]

A study showed the potential relationship between FI and life satisfaction (LS).^[13] LS is an important component of well-being, indicating the degree to which people can effectively cope with various changes and conditions.^[14] LS improves positive feelings and motivation for engagement in physical activities. Hence, individuals with higher LS have better functional abilities and use better strategies to deal with problems.^[14] LS is also correlated with physical, mental, and social factors,^[15] FI, social life, income, education, mental health, and satisfaction with peers.^[15,16]

A study reported that improvement in FI through regular physical and leisure activities can improve LS among older adults.^[17] A study found that compared with the time spent on ADL and rest, the time spent on leisure and occupational activities was associated with higher levels of LS.^[18] Another study reported that older adults without impairment in performing ADL had higher levels of LS.^[19] We can suppose that COVID-19 and its consequences may cause functional limitations in older adults and affect their FI and LS, but, contradictory results can be found. For example, a Chinese study showed that the majority of participants reported LS despite prolonged isolation.^[20] Despite the importance of FI and LS among older adults, there are limited data about this relationship in this population in Iran. The restrictions caused by the COVID-19 condition, including long-term quarantine and social distancing, affect the lives of older adults. Therefore, the present study was conducted to produce more evidence in this area.

Objectives

This study aimed to evaluate the relationship between FI and LS among older adults during the COVID-19 pandemic.

Methods

Study Design and Participants

This cross-sectional study was conducted in 2021. The study population consisted of all community-dwelling older adults who were referred to the public healthcare centers in southern Tehran, Iran. All these centers are covered by Tehran University of Medical Sciences. Cluster sampling was performed to select the health centers. Each of the five districts in southern Tehran was considered a cluster from which two or three public healthcare centers were randomly selected. An equal number of eligible older adults were then selected from each center through simple random sampling. Eligibility criteria were age 60 years and older, basic literacy skills, dwelling in the community, and no debilitating chronic diseases or cognitive or mental problems (as self-reported by the participants and chart review). The sample size was calculated with a confidence level of 0.95, a power of 0.90, and a LS-FI correlation coefficient of at least 0.15 to be considered statistically significant.^[21] The formula for calculating sample size [Formula 1] revealed that at least 479 participants were needed.

$$\frac{(Z_{1-\alpha/2} + Z_{1-\beta})^2}{W^2} + 3, \quad W = 0.5 \times \ln[(1+r)/(1-r)]$$

Formula 1. Sample size calculation formula

Data collection instruments

The research questionnaires were completed by the subjects with the help of the researcher in the first meeting. However, due to the conditions of COVID-19, some information was completed over the phone. In the health centers, all the older adults in the region are registered in the Sib system, and some demographic information could be obtained from the Sib system.

Data were collected using a demographic questionnaire, the ADL scale, the IADL scale, and the Life Satisfaction Index-Z. The items of the demographic questionnaire were age, gender, weight, height, marital status, employment status, education level, income, number of children, and medications used.

We used the eight-item ADL scale and the seven-item IADL scale. The items of both scales are scored on a three-point scale as follows: “zero: Dependent”, “1: Needs help”, and “2: Independent”. The total scores for the eight-item ADL and the seven-item IADL scales are respectively 0–16 and 0–14 which are interpreted as follows: scores 0–7 for ADL and 0–6 for IADL: dependent; scores 8–11 for ADL and 7–10 for IADL: needs help; and scores 12–16 for ADL and 11–14 for IADL: independent.^[7]

In a previous study in Iran, Cronbach's alpha and the test-retest intraclass correlation coefficient were reported to be 0.80 and 0.76 for the ADL scale and 0.75 and 0.79 for the IADL scale.^[22]

The Life Satisfaction Index-Z has five negatively-worded items (i.e., items 3, 6, 10, 11, and 13) and eight positively worded items (i.e., items 1, 2, 4, 5, 7, 8, 19, and 12). Positively worded items are scored "zero: I don't know", "1: Disagree", or "2: Agree" and negatively worded items are scored reversely. Accordingly, the total possible score of the index is 0–26, which is interpreted as follows: 0–12: low LS, 13–21: moderate LS, and 22–26: high LS. A study of older adults in Iran evaluated the psychometric properties of this index and reported that its test-retest intraclass correlation coefficient, Cronbach's alpha, and split-half unequal length Spearman-Brown coefficient were 0.93, 0.79, and 0.79, respectively.^[23]

Data analysis

The SPSS software (v. 16.0) was employed (SPSS Inc., Chicago, IL, USA) for data analysis at a significance level of <0.05. Descriptive statistics (frequency, percent, mean, and standard deviation) were used to describe participants' characteristics. The independent sample t-test and analysis of variance were used to compare the mean scores of LS between subgroups of participants. Pearson's correlation coefficient was used to examine the correlation between dependent and independent variables. Multiple linear regression analysis with the Enter method was used to determine the variables associated with LS. All variables with $P \leq 0.2$ in univariate analysis were entered into the model as independent variables.

Ethical considerations

This study was performed by observing the Declaration of Helsinki. The Ethics Committee of Tehran University of Medical Sciences, Tehran, Iran, approved this study (code: IR.TUMS.MEDICINE.REC.1399.1118). Participants were provided with information about the study aim and the confidential management of their information and their informed consent was obtained. Due to the pandemic conditions, we tried to implement all health protocols.

Results

A total of 479 older adults participated in this study. Most participants were male (54.1%), married (71.8%), and retired (52%), 44.3% of them had primary education [Table 1].

The mean score of LS was 11.94 ± 5.11 and 54.9% of participants had low LS, while only 2.5% of them had high LS. The mean scores of FI in ADL and IADL were also

15 ± 2.32 and 11.28 ± 3.72 and the level of FI in ADL and IADL was 90.4% and 73.3%, respectively [Table 2].

LS had a significant relationship with education level and income ($P \leq 0.001$), FA in ADL had a significant relationship with gender, marital status, and employment status ($P < 0.001$), and FI in IADL had a significant relationship with gender, marital status, employment status, education level, and income ($P < 0.001$) [Table 1]. Moreover, LS had a significant correlation with FI in ADL ($r = 0.133$; $P = 0.004$) and IADL ($r = 0.213$; $P < 0.001$).

Regression analysis revealed that FI in IADL ($P = 0.013$) and income ($P < 0.001$) were significantly associated with LS. Accordingly, each one-point increase in the score of FI in IADL was associated with a 0.241 point increase in the score of LS. Moreover, the LS of participants with insufficient income and those with relatively sufficient income were 5.77 and 3.499 points lower than those with sufficient income, respectively. Independence in instrumental ADL and income significantly predicted 30% of the variance of LS ($P < 0.05$, [Table 3]).

Discussion

Findings revealed a significant positive relationship between ADL and IADL with LS. The majority of participants were independent in ADL and IADL. Studies show that disabilities increase significantly with age.^[24] A study in Turkey showed that FI was clearly lower in the age group over 80 than those in the 65-70 age group.^[5] A study of 300 older adults in Iran also reported that 55% of them were completely independent in ADL and 90% were relatively independent in IADL.^[25] Another study of older adults in rural areas in Iran also reported that they were relatively independent.^[26] The high level of independence in the present study is attributable to the fact that the study was conducted during the COVID-19 pandemic and older adults had to perform their ADL and IADL as independently as possible due to the need for quarantine and physical distancing. The results of studies on the effects of the COVID-19 pandemic on older adults' FI are inconsistent. Some studies have shown that quarantine and physical distancing could decrease physical activity and increase dependence among older adults.^[27,28] However, a study found no significant difference between the mean scores of FI in ADL and IDAL before and ninety days after affliction by COVID-19 among Indian older adults.^[29] These inconsistent results are attributable to differences among studies respecting their participants' characteristics such as age, place of residence, and geographic area, and highlight the importance of further studies in this area.

Table 1. Participants' demographic characteristics and their relationship with life satisfaction and functional independence

Characteristics	n (%) or Mean±SD (Range)	Life satisfaction		Activities of daily living		Instrumental activities of daily living	
		Mean±SD	P value	Mean±SD	P value	Mean±SD	P value
Gender							
Female	220 (45.9)	12.17 (4.94)	0.372 ^a	14.66±2.59	0.004	10.54±4.10	< 0.001
Male	259 (54.1)	11.75 (5.26)		15.28±2.02		11.91±3.24	
Marital status							
Married	344 (71.8)	12.21 (5.02)	0.063 ^a	15.42±1.71	<0.001	12.15±2.85	< 0.001
Single/divorce d/widowed	135 (28.2)	11.25 (5.3)		13.91±3.17		9.05±4.65	
Employment status							
Retired	249 (52)	12.16 (5.19)	0.55 ^a	14.99±2.30	<0.001	11.17±3.81	< 0.001
Housewife	158 (33)	11.82 (4.84)		14.62±2.71		10.55±4.01	
Self-employed	72 (15)	11.45 (5.46)		15.86±1.58		13.27±2.28	
Education level							
Primary	212 (44.3)	10.66 (4.86)	< 0.001 ^b	14.72±2.67	0.058	10.39±4.15	< 0.001
Guidance school	66 (13.8)	12.1 (4.31)		15.19±2.09		11.43±3.07	
Diploma	106 (22.1)	12.75 (4.83)		15.01±2.34		11.62±3.66	
University	95 (19.8)	13.8 (5.75)		15.47±1.34		12.78±2.47	
Income							
Insufficient	136 (28.4)	9.01 (4.06)	< 0.001 ^b	14.72±2.43	0.114	10.94±3.91	0.009
Relatively sufficient	190 (39.7)	11.33 (4.59)		14.96±2.45		10.92±3.87	
Sufficient	153 (31.9)	15.31 (4.68)		15.28±2.02		12.04±3.24	
Age (Years)	72.82±7.33 (60–97)	r=-0.039 P=0.394 ^c		r=-0.499 P<0.001 ^c		r=-0.567 P<0.001 ^c	
Body mass index	26.18±3.72 (17.1–45.2)	r=-0.02 P=0.656 ^c		r=-0.067 P=0.145 ^c		r=-0.045 P=0.328 ^c	

^a The results of the independent-sample t-test, ^b The results of the one-way analysis of variance, ^c The results of Pearson's correlation analysis.

Table 2. The mean scores and levels of life satisfaction and functional independence

Variables	n (%)	Mean±SD (Range)
Activities of daily living		15±2.32 (4–16)
Dependent	15 (3.1)	
Needs help	31 (6.5)	
Independent	433 (90.4)	
Instrumental activities of daily living		11.28±3.72 (0–14)
Dependent	64 (13.4)	
Needs help	64 (13.4)	
Independent	351 (73.2)	
Life satisfaction		11.94±5.11 (1–25)
Low	263 (54.9)	
Moderate	204 (42.6)	
High	12 (2.5)	

Table 3. Results of multiple linear regression analysis for the prediction of life satisfaction based on functional independence and demographic characteristics

Independent variables	Beta		T	P value	Confidence interval	R ²
	Non-standardized	Standardized				
Independence in activities of daily living	-0.235	-0.107	-1.579	0.115	-0.528, 0.057	0.3
Instrumental independence in activities of daily living	0.241	0.175	2.492	0.013	0.051, 0.431	
Marital status						
Married	-0.174	-0.015	-0.355	0.723	-1.136, 0.788	
Single (Reference)						
Education level						
Primary	-1.076	-0.104	-1.853	0.065	-2.216, 0.065	
Guidance school	-0.030	-0.002	-0.042	0.966	-1.444, 1.383	
Diploma	-0.123	-0.010	-0.197	0.844	-1.357, 1.110	
University (Reference)						
Income						
Insufficient	-5.776	-0.509	-10.857	0.000	-6.821, -4.730	
Relatively sufficient	-3.499	-0.335	-7.152	0.000	-4.461, -2.538	
Sufficient (Reference)						

Our findings also indicated a significant positive relationship between LS and FI in ADL and IADL. Independence is a key factor in the fulfillment of basic needs, and individuals that are more independent have higher LS. Like our findings, a study showed that physical activity was positively associated with LS among Spanish older adults.^[30] Some other studies also reported that functional limitations in ADL and IADL reduce social interactions and LS among older adults.^[13,31] Contradictorily, a study reported a significant inverse relationship between LS and FA in IADL.^[32] Another study also showed that older adults had higher LS compared to their younger counterparts because they were less responsive to negative conditions.^[15] This discrepancy might be related to the difference in education level of participants in different studies, because education level is significantly related to LS^[33] and FI^[34] among older adults. Engaging in most activities needs adequate knowledge and skills, hence, older adults with higher education levels can manage their activities better and thereby, feel higher levels of LS.

Study findings also revealed FI in IADL and income as the significant predictors of LS among older adults. The effect of income on LS was also stronger than that of other variables in the regression model. Consistent with our findings, a study showed that financial status had a significant positive relationship with LS.^[35] Other studies also confirmed the role of financial status in LS and quality of life of older adults.^[2,32]

One of the most important limitations of this study was the fourth and the fifth waves of the COVID-19 pandemic in Iran during sampling and data collection, so we had to collect some data through telephone contact. Moreover,

participants' psychological status during data collection might have affected the results.

Conclusions

Our older adults had optimal levels of FI in ADL and IADL and a low level of LS. FI in ADL and IADL has a significant relationship with LS, while the significant predictors of LS are FI in IADL and income. Healthcare authorities can take steps to promote the FI of older adults and thereby improve their LS, by carrying out interventions such as formulating health policies, amending health system laws, and establishing insurance coverage to reduce costs. Furthermore, by addressing various dimensions of older adults' health, especially during epidemics such as the COVID -19 pandemic, community health nurses can also improve the LS of older adults. Policymakers and health planners should also consider the necessary measures to regularly monitor FI and LS of older adults. It is suggested that similar studies be conducted with more samples, and in other settings and under non-pandemic conditions.

Acknowledgment

We would like to thank the Student Research Center of Tehran University of Medical Sciences, Tehran, Iran, for financially supporting this study. Moreover, we are thankful to all participants of the study as well as the staff of the study setting.

Competing interests

None.

Abbreviations

functional independence: FI
 life satisfaction: LS
 activities of daily living: ADL
 Instrumental ADL: IADL

Authors' contributions

All authors read and approved the final manuscript. All authors take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This study was funded and supported by the Tehran University of Medical Sciences (TUMS); (Grant: 50551).

Role of the funding source

None.

Availability of data and materials

The data used in this study are available from the corresponding author on request.

Ethics approval and consent to participate

This study was performed by observing the Declaration of Helsinki. The Ethics Committee of Tehran University of Medical Sciences, Tehran, Iran, approved this study (code: IR.TUMS.MEDICINE.REC.1399.1118). Participants were provided with information about the study aim and the confidential management of their information and their informed consent was obtained. In addition, due to the pandemic conditions, we tried to implement all health protocols.

Consent for publication

By submitting this document, the authors declare their consent for the final accepted version of the manuscript to be considered for publication.

References

1. Esmaili M. The effect of physical activities on the quality of life, hope and life satisfaction among the elderly in Ilam city. *J Gerontol* 2017;2:29-40. doi:10.29252/joge.2.1.29
2. Abolhasani F, Bastani F. Successful Ageing in the Dimensions of Life Satisfaction and Perception of Ageing in the Iranian Elderly Adults Referring to the Health Center in the West of Tehran, Iran. *Iran J Nurs* 2019;31:61-74. doi:10.29252/ijn.31.116.61
3. Fahimfar N, Noorali S, Yousefi S, Gharibzadeh S, Shafiee G, Panahi N, *et al.* Prevalence of osteoporosis among the elderly population of Iran. *Arch Osteoporos* 2021;16:1-10. doi:10.1007/s11657-020-00872-8 PMID:33475880
4. Koç Z. The investigation of factors that influence self-care agency and daily life activities among the elderly in the northern region of Turkey. *Collegian* 2015;22:251-8. doi:10.1016/j.colegn.2014.01.002 PMID:26552195
5. Bozkurt Ü, Yılmaz M. The determination of functional independence and quality of life of older adults in a nursing home. *Int J Caring Sci* 2016;9:198-210.
6. Sadegh Moghadam L, Foroughan M, Mohammadi F, Ahmadi F, Farhadi A, Nazari S, *et al.* Aging perception in older adults. *Iran J Ageing* 2016;10:202-9.
7. Mortazavi H, Tabatabaiechehr M, Taherpour M, Masoumi M. Investigating the Status of Daily Life Activities (Basic, Instrumental, Advanced) and Related Factors in the Elderly. *J North Khorasan Univ Med Sci* 2020;12:88-95. doi:10.52547/nkums.12.2.88
8. Safa A, Alavi N, Abedzadeh-Kalahroudi M. Predictive factors of dependency in activities of daily living following limb trauma in the elderly. *Trauma monthly* 2016;21:e25091. doi:10.5812/traumamon.25091 PMID:28184359 PMCid:PMC5292017
9. Ishida A, Ishida E. Changes in Daily Life Satisfaction among Community Dwelling Elderly during the COVID-19 Pandemic in Japan. *J Ageing Longevity* 2021;1:3-10. doi:10.3390/jal1010002
10. Tran B, Allnutt A, Wong A. COVID-19 pandemic and youth fitness: a systematic review. *Novel Clin Med* 2023;2:3-10. doi:10.22034/ncm.2023.381863.1069
11. Emily L, Gilad G, Haim ME, Galina G. Functional dependency and COVID-19 in elderly patients with mild to moderate disease. Experience of tertiary geriatric hospital. *Exp Gerontol* 2022;157:111620. doi:10.1016/j.exger.2021.111620 PMID:34742855 PMCid:PMC8564951
12. Pashaeypoor S, Baumann SL, Sadat Hoseini A, Cheraghi MA, Chenari HA. Identifying and Overcoming Barriers for Implementing Watson's Human Caring Science. *Nurs Sci Q* 2019;32:239-44. doi:10.1177/0894318419845396 PMID:31203774
13. Chauhan S, Kumar P, Srivast S, Patel R. Do Functional Limitations Predict Life Satisfaction Among Older Adults in India: A Study based on LASI Survey in India. *Res Sq* 2021:1-22. doi:10.21203/rs.3.rs-721491/v1
14. Jamali Moghaddam M, Bastani F. Investigate Life Satisfaction and Correlates in Older Adults Attending West Health Center of Tehran, 2017. *Iran J Nurs Res* 2019;14:70-80.
15. Pan Y, Chan SHW, Xu Y, Yeung KC. Determinants of life satisfaction and self-perception of ageing among elderly people in China: An exploratory study in comparison between physical and social functioning. *Arch Gerontol Geriatr* 2019;84:103910. doi:10.1016/j.archger.2019.103910 PMID:31302503
16. Khodabakhsh S. Factors Affecting Life Satisfaction of Older Adults in Asia: A Systematic Review. *J Happiness Studies* 2022;23:1289-304. doi:10.1007/s10902-021-00433-x
17. Liu LH, Kao CC, Ying JC. Functional capacity and life satisfaction in older adult residents living in long-term care facilities: The mediator of autonomy. *J Nurs Res* 2020;28:e102. doi:10.1097/JNR.0000000000000362 PMID:31904735
18. Maher J, Pincus AL, Ram N, Conroy DE. Daily physical activity and life satisfaction across adulthood. *Dev Psychol* 2015;51:1407-1419. doi:10.1037/dev0000037 PMID:26280838 PMCid:PMC4579061
19. Boccaccio DE, Cenze I, Covinsky KE. Life satisfaction among older adults with impairment in activities of daily living. *Age Ageing* 2021;50:2047-54. doi:10.1093/ageing/afab172

- PMid:34510173 PMCID:PMC8581387
20. Wang X, Lei SM, Le S, Yang Y, Zhang B, Yao W, *et al.* Bidirectional Influence of the COVID-19 Pandemic Lockdowns on Health Behaviors and Quality of Life among Chinese Adults. *Int J Environ Res Public Health* 2020;17:5575. doi:10.3390/ijerph17155575 PMid:32748825 PMCID:PMC7432516
 21. Hulley S. Designing clinical research. Third ed: Lippincott Williams & Wilkins; 2007.
 22. Taheri Tanjani P, Azadbakht M. Psychometric properties of the Persian version of the activities of daily living scale and instrumental activities of daily living scale in elderly. *J Mazandaran Univ Med Sci* 2016;25:103-12.
 23. Tagharrobi Z, Tagharrobi L, Sharifi K, Sooki Z, Nele S, Ghotbi N, *et al.* Psychometric evaluation of the Life Satisfaction Index-Z (LSI-Z) in an Iranian elderly sample. *Health Monit* 2011;10:5-13.9.
 24. Tavafian S, Aghamolaei T, Moeini B. Functional independence level of physical activities in elderly people: a population-based study. *Health Monitor* 2014;13:449-56.
 25. Nourbakhsh S, Fadayevatan R, Alizadeh-Khoei M, Sharifi F. Determining the status of activity of daily living (ADL) and instrumental activity of daily living (IADL) in healthy and cognitive impaired elderlies. *Jorjani Biomed J* 2017;5:63-77.
 26. Jokar F, Asadollahi AR, Kaveh MH, Ghahramani L, Nazari M. Relationship of Perceived Social Support With the Activities of Daily Living in Older Adults Living in Rural Communities in Iran. *Iran J Ageing* 2020;15:350-65. doi:10.32598/sija.10.15.3.2773.2
 27. Lebrasseur A, Fortin-Bédard N, Lettre J, Raymond E, Bussi eres EL, Lapi erie N, *et al.* Impact of the COVID-19 pandemic on older adults: rapid review. *JMIR aging* 2021;4:e26474. doi:10.2196/26474 PMid:33720839 PMCID:PMC8043147
 28. Yamada M, Kimura, Y, Ishiyama D, Otobe Y, Suzuki M, Koyama S, *et al.* Effect of the COVID-19 Epidemic on Physical Activity in Community-Dwelling Older Adults in Japan: A Cross-Sectional Online Survey. *J Nutr Health Aging* 2020;24:948-50. doi:10.1007/s12603-020-1501-6 PMid:33155619 PMCID:PMC7597428
 29. P S, Madhavan S, Pandurangan V. Prevalence, Pattern and Functional Outcome of Post COVID-19 Syndrome in Older Adults. *Cureus* 2021;13:e17189.
 30. Parra-Rizo MA, Sanchis-Soler G. Satisfaction with life, subjective well-being and functional skills in active older adults based on their level of physical activity practice. *Int J Environ Res Publ Health* 2020;17:1299. doi:10.3390/ijerph17041299 PMid:32085450 PMCID:PMC7068550
 31. Steptoe A, Di Gessa G. Mental health and social interactions of older people with physical disabilities in England during the COVID-19 pandemic: a longitudinal cohort study. *Lancet Public health* 2021;6:e365-e373. doi:10.1016/S2468-2667(21)00069-4 PMid:33894138
 32. Ng ST, Tey NP, Asadullah MN. What matters for life satisfaction among the oldest-old? Evidence from China. *PLoS one* 2017;12:e0171799. doi:10.1371/journal.pone.0171799 PMid:28187153 PMCID:PMC5302476
 33. Yang DC, Lee JD, Huang CC, Shih HI, Chang CM. Association between multiple geriatric syndromes and life satisfaction in community-dwelling older adults: A nationwide study in Taiwan. *Arch Gerontol Geriatr* 2015;60:437-42. doi:10.1016/j.archger.2015.02.001 PMid:25726424
 34. Moeini B, Barati M, Jalilian F. Factors associated with the functional independence level in older adults. *Hormozgan Med J* 2012;15:318-26.
 35. Hsu HC. Trajectory of life satisfaction and its relationship with subjective economic status and successful aging. *Soc Indicat Res* 2010;99:455-68. doi:10.1007/s11205-010-9593-8.

How to Cite this Article:

Saghaei MH, Salehpour-Emran M, Sarouei N, Zarekar M, Haghani SH, Pashaeypoor SH. The relationship between functional independence and life satisfaction among Iranian community-dwelling older adults during the COVID-19 pandemic. *Nurs Midwifery Stud.* 2023;12(2):158-164. doi:10.48307/NMS.2023.179817